



Australian Government
Australian Sports Commission



CONCUSSION AND BRAIN HEALTH POSITION STATEMENT 2024

An initiative of the Australian Institute of Sport,
Sports Medicine Australia, Australian Physiotherapy Association
and Australasian College of Sport and Exercise Physicians

June 2024

“If in doubt, sit them out”



AUSTRALASIAN COLLEGE OF
SPORT AND EXERCISE PHYSICIANS



AUSTRALIAN
PHYSIOTHERAPY
ASSOCIATION



SPORTS
MEDICINE
AUSTRALIA

CONTENTS

GLOSSARY	4
EXECUTIVE SUMMARY	6
INTRODUCTION	7
Aims	7
MEDICO-LEGAL CONSIDERATIONS	8
WHAT IS CONCUSSION?	8
CONCUSSION EPIDEMIOLOGY	9
Concussion epidemiology in high performance sports	9
Sex differences in concussion	9
Concussion epidemiology in high-performance para-athletes	10
Concussion epidemiology in recreational sports	12
Concussion epidemiology in youth sports	13
Economic impact of concussion	13
PATHOPHYSIOLOGY	14
RISK REDUCTION/PREVENTION	15
RECOGNISE, REMOVE, REFER	16
Clinical features of concussion	17
Mandatory signs of concussion (immediate removal and no further play)	17
Discretionary signs of concussion (immediate removal for further assessment)	18
Red flag signs of concussion (immediate referral to emergency department)	18
RE-EVALUATE	19
Symptoms of concussion	19
Diagnosis of concussion	20
Concussion assessment in para-athletes	21
Concussion assessment in the community	22
Health care practitioner concussion evaluation in a controlled office environment	22
Concussion assessment in children aged 5–12 years	23
Health care practitioner concussion evaluation in children in a controlled office environment	25
CONCUSSION MANAGEMENT	30
The role of physiotherapy in concussion	30
Interdisciplinary care for athletes with concussion	30
RELATIVE REST	32

REHABILITATION	32
Vestibular and oculomotor impairments	32
RETURN TO SPORT FOR COMMUNITY AND YOUTH SETTINGS	34
Concussion officer for community/school environments	36
ADVANCED CARE RETURN TO SPORT FRAMEWORK	38
Protocol for those with multiple suspected concussions	38
EMERGING TOOLS FOR CONCUSSION DIAGNOSIS	40
Emerging tools	40
Vestibular and oculomotor assessment	40
CONCUSSION AND LONG-TERM BRAIN HEALTH	41
Long-term brain health in female athletes	43
Long-term brain health in para-athletes/athletes with disabilities	43
HOLISTIC BRAIN HEALTH AND WELLBEING IN CURRENT AND FORMER ATHLETES	44
QUESTIONS TO BE ANSWERED IN IMPROVING UNDERSTANDING OF CONCUSSION AND LONG-TERM BRAIN HEALTH	45
EDUCATION	46
Education modules	46
KEY POINTS FOR ATHLETES, COACHES, PARENTS, TEACHERS AND SUPPORT STAFF	47
KEY POINTS FOR HEALTHCARE PRACTITIONERS	48
REFERENCES	49
APPENDIX A — PHYSIOTHERAPY GUIDED REHABILITATION OF CONCUSSION	60

GLOSSARY

AAN	American Academy of Neurology
ACRM	The American Congress of Rehabilitation Medicine [ACRM]
ACS	Advanced Care Setting
ADHD	Attention Deficit Hyperactivity Disorder
AFL	Australian Football League
AIS	Australian Institute of Sport
ASC	Australian Sports Commission
AUDIT	Alcohol Use Disorders Identification Test
BESS	Balance Error Scoring System
CBHPS24	Concussion and Brain Health Position Statement 2024
Child-SCAT6	Child Sport Concussion Assessment Tool 6
Child-SCoAT6	Child Sport Concussion Office Assessment Tool-6
CIPS	Concussion in Para Sport
CISG	Concussion in Sport Group
CMT	Clinical Management Team (a group of healthcare practitioners, ideally including a medical practitioner, physiotherapist, and, depending on the complexity of the case, possibly other healthcare professional such as a psychologist, neuropsychologist, neurologist etc.)
CP	Cerebral Palsy
CRT6	Concussion Recognition Tool 6
CTE	Chronic Traumatic Encephalopathy
CTE-NC	Chronic Traumatic Encephalopathy Neuropathological Change
DASS	Depression, Anxiety, and Stress Scale
fMRI	Functional Magnetic Resonance Imaging
GRTS	Graded Return to Sport
GRTSF	Graded Return to Sport Framework
HBI	Health and Behaviour Inventory
HCP	Healthcare Practitioner (AHPRA registered health care practitioner with appropriate training and experience in concussion assessment and management)
IOC	International Olympic Committee
IPC	International Paralympic Committee
IQR	Interquartile Range
MILS	Manual in-line stabilisation
MPCS	Melbourne Paediatric Concussion Scale
MRS	Magnetic Resonance Spectroscopy
mTBI	Mild Traumatic Brain Injury
NCAA	National Collegiate Athletic Association
NIN	National Institute Network

NPC	Near Point of Convergence
NRL	National Rugby League
NSO	National Sporting Organisation
NSOD	National Sporting Organisations for People with Disability
PCS	Post-Concussive Syndrome
PCSI	Post Concussion Symptom Inventory
PCSQ	Post-Concussive Syndrome Questionnaire
PET	Positron Emission Tomography
PoC	Point of Care
PPE	Personal protection equipment
p-tau	Hyperphosphorylated Tau
RHT	Repeated Head Trauma
SCAT6	Sport Concussion Assessment Tool 6
SCI	Spinal Cord Injury
SCOAT6	Sport Concussion Office Assessment Tool-6
sncRNAs	Salivary Small Non-Coding RNAs
TES	Traumatic Encephalopathy Syndrome
US CDC	The United States Centers for Disease Control and Prevention
VOM	Vestibular and Oculomotor
VOMS	Vestibular and Oculomotor System

EXECUTIVE SUMMARY

There has been growing concern in sporting communities in Australia and internationally about the potential health ramifications for athletes from repeated head trauma (RHT) and sport-related concussion (referred to as *concussion* here on). Concussion affects athletes at all levels of sport, from the part-time recreational athlete to the full-time professional. If managed appropriately, most episodes of concussion resolve over a short period of time, with or without medical intervention. Complications can occur, however, including prolonged duration of symptoms and increased susceptibility to further injury. There is also concern about potential consequences of RHT for long term brain health.

Chronic traumatic encephalopathy (CTE) is a neurodegenerative pathology associated with a history of RHT. There are a growing number of case studies and case series which document CTE neuropathological change (CTE-NC) in retired athletes with a history of RHT. There remain many uncertainties about the strength of the association between RHT, concussion, and CTE-NC. Further research is required to understand the prevalence of CTE-NC in athletic cohorts and the factors that predispose some athletes to the development of CTE-NC following exposure to RHT.

Over recent years there has been elevated public awareness of concussion and increased focus on the importance of diagnosing and managing the condition promptly, safely, and appropriately.

Sport administrators, healthcare practitioners (HCPs), coaches, parents/guardians/caregivers, and athletes are seeking information regarding the timely recognition and appropriate management of concussion. There is a need for clear, unequivocal, and reliable information to be readily accessible to all members of the community. There is a need for clarity of message and consistency of message in order to optimise safety in contact, collision and combat sports.

The Australian Sports Commission (ASC) is the Australian Government agency responsible for supporting and investing in sport at all levels. The ASC's strategic vision is to ensure sport has a place for everyone and delivers results that make Australia proud. The ASC also plays a critical leadership role in guiding sporting organisations and the sport sector in relation to a range of issues impacting sport. The ASC is not a regulatory authority and has no power to enforce compliance or regulations. The Australian Institute of Sport (AIS) is the high-performance arm of the ASC. As the high-performance arm of the ASC, the AIS functions as a resource for sport organisations, providing expertise and education as required. The AIS seeks to guide, but not instruct sports on a range of policy positions, including concussion. Funded by the Australian Government, this Concussion and Brain Health Position Statement 2024 brings together the most contemporary evidence-based information and presents it in a format that is appropriate for all stakeholders. This position statement is an update to the previous position statement, launched in 2023. The 2024 update is intended to ensure that participant safety and welfare is paramount when dealing with matters of RHT and concussion in sport.

This update includes the latest recommendations and addresses evidence specifically relating to the management and treatment of concussion by physiotherapists as healthcare practitioners. This position statement provides an overview of the scientific literature in relation to the potential long-term effects of RHT and concussion in athletes. It recommends that sports operate on a principle of an 'abundance of caution'. **'If in doubt, sit them out'**. Where there is any suspicion of concussion, an athlete should be removed from the field of play and should not be allowed to return, until the successful completion of a graduated return to sport/learn program, and clearance by a healthcare practitioner.

This position statement is intended as a general guide relating to current best practices for prevention, recognition, and management of concussion, underpinned by available empirical evidence. It should not be interpreted as a policy for clinical practice or legal standard of care. Recommendations will evolve to reflect evidence and advances in science.

INTRODUCTION

Sport-related concussion [referred to as *concussion* here on] is a public health issue of concern, in Australia and globally. Repeated head trauma (RHT) includes head impacts that lead to a concussion or a mild traumatic brain injury, as well as head trauma that *do not* cause an individual to experience any subsequent symptoms. Contact or collisions during sport participation is one way an individual may experience RHT. 16% of all annual concussions that required hospital treatment in Victoria are related to participation in sport.¹

Concussion affects male and female athletes, para and able bodied, across all age groups and at all levels of sport. The risk of concussion can vary across sports, level of participation, session type (i.e., training or competition), and sex.² The rate of concussion has been reported to be higher in females than in males.^{2,3} Individuals who participate in contact, collision, and combat sports have a high risk of RHT and concussion as they can be exposed to multiple blows to the head (direct) and trunk (indirect).² It is important to note that although blows to the trunk may not directly impact the head, it can also cause concussion. As contact and collision sports gain popularity among groups such as females, and athletes with disabilities, it is anticipated that a concurrent rise in concussion injuries may be observed in these groups.

There is increasing concern about the long-term effects on the brains of individuals who experience RHT or multiple concussion, driven in part by highly publicised cases of dementia and mental health issues among American football players, boxers, rugby players, and Australian Football League (AFL) players. Concern about the possible short-term and long-term health ramifications for athletes has led to an increased focus on improving prevention, recognition, and management of concussion.

The Australian Sports Commission (ASC) is the Australian Government agency responsible for supporting and investing in sport at all levels. The ASC's strategic vision is to ensure sport has a place for everyone and delivers results that make Australia proud. The ASC also plays a critical leadership role in guiding sporting organisations and the sport sector in relation to a range of issues impacting sport. The ASC is not a regulatory authority and has no power to enforce compliance or regulations. The Australian Institute of Sport (AIS) is the high-performance arm of the ASC. The AIS functions as a resource for sport organisations, providing expertise and education as required. The AIS seeks to guide, but not instruct sports on a range of policy positions, including concussion.

Aims

There has been a rapid increase of publications on concussion. For instance, the United States National Library of Medicine's PubMed database had 1,840 concussion-related publications in 2021, 1,600 in 2022 and a further 1,340 in 2023. The AIS is committed to ensure that everyone involved in sport including athletes, coaches, parents/guardians/caregivers, health care practitioners, and others can access the latest evidence-based resources on concussion. Funded by the Australian Government, the Concussion and Brain Health Position Statement 2024 (CBHPS24) aims to:

- > provide access to up-to-date evidence-based information on concussion for all Australians
- > provide improved safety and health outcomes for all individuals who suffer concussive injuries while participating in sport
- > assist Australian sporting organisations to align their policy and procedures to the most up-to-date scientific evidence
- > protect the integrity of sport through consistent application of best practice protocols and guidelines
- > provide a platform to support the development of a national policy for the management of concussions in Australia
- > Provide clarity of message and consistency of message in articulating recommendations to stakeholders in sport.

It should be noted that this position statement operates on a principle of an abundance of caution [**'if in doubt, sit them out'**]. Where there is any suspicion of concussion, an athlete should be removed from the field of play and should not be allowed to return, until the successful completion of a graduated return to sport/learn program, and clearance by a healthcare practitioner. The recommended return to sport/learn protocol is dependent on the level of healthcare cover available, the age of the athlete and the fulfilment of eligibility criteria to have a more advanced care protocol established. The [Concussion in Sport Australia](#) website hosts freely available, evidence-based resources to accompany the CBHPS24 for better knowledge translation to improve athlete care at all levels of sport.

MEDICO-LEGAL CONSIDERATIONS

The CBHPS24 is intended as a general guide relating to current best practices for prevention, recognition, and management of concussion, underpinned by available empirical evidence. It will be updated as new knowledge develops. There is recognition of nuanced differences across sport settings, rules, resources, and culture, between sports, and from recreational to high performance/professional sports. The information conveyed in this position statement does not constitute medical, legal, or other professional advice; This position statement therefore **should not** be interpreted as a guideline for clinical practice or legal standard of care. Recommendations will evolve to reflect evidence and advances in science. Owing to differences in risk profiles, rules, settings and resources, concussion guidelines need to be adopted for individual sport specific regulatory environments. Also, given the complex nature of recognition and management of concussion, individual treatments may depend on the circumstances of each case. At the same time, consistent application of the principles highlighted in this position statement and International Olympic Committee (IOC) consensus statement guidelines can facilitate a greater degree of standardisation in the recognition and management of concussion.

The CBHPS24 is viewed as a 'live' document and will be updated as research and evidence become available in this rapidly evolving space of sport.

WHAT IS CONCUSSION?

A concussion occurs through a collision with another person or object where biomechanical forces to the head, or anywhere on the body transmit an impulsive force to the head/brain. In most concussions, this results in transient neurological impairment. It should be noted that concussion can also occur with relatively minor 'knocks'. Concussion is often an evolving injury with symptoms changing over hours or days following the injury.^{4,5} There are often adverse effects on balance and cognitive function.⁴ Recovery times following concussion vary between athletes. Pathophysiological recovery may take longer than measures of clinical recovery,⁶ and the average time taken to resolve clinical symptoms may vary according to sex, age, presence of pre-injury medical conditions, and para-athlete status.^{3,7-9} The current clinical definition of concussion does not distinguish persistent symptoms, or the underlying processes that impair brain function or any potential brain abnormalities. To overcome this limitation, the Amsterdam panel of the Concussion in Sport Group (CISG) defined concussion as *"a traumatic brain injury caused by a direct blow to the head, neck or body resulting in an impulsive force being transmitted to the brain that occurs in sports and exercise-related activities. ... Symptoms and signs may present immediately, or evolve over minutes or hours, and commonly resolve within days, but may be prolonged."*¹⁰

Most concussions are limited to physiological changes with no identifiable structural change. In some cases, there may be microscopic structural changes which would position those cases within the traumatic brain injury spectrum. As discussed below, some epidemiological data, particularly hospital data, do not distinguish between traumatic brain injury and concussion.

The American Congress of Rehabilitation Medicine Diagnostic Criteria (ACRM) consider a concussion to be a mild traumatic brain injury (mTBI). A mTBI is 'suspected' if an athlete experiences symptoms that are believed to arise from a concussion, but there are no clinical signs and no objective clinical examination findings.¹¹ It should be noted that both the CISG definition and ACRM diagnostic criteria indicate that an athlete who develops symptoms consistent with concussion should be removed from play and undergo a graded return to sport strategy^{10, 11} and align with '**when in doubt, sit them out**' mantra.

CONCUSSION EPIDEMIOLOGY

Currently, Australia does not have a national sports injury surveillance system. Therefore, precise data on the incidence, frequency, and prevalence of concussion in Australia is undetermined. This is further compounded by a lack of recognition of the signs and symptoms of concussion,^{12, 13} under-reporting and failing to seek medical advice.¹⁴⁻¹⁶

Furthermore, the terms 'sex' (biology) and 'gender' (social construct) are inconsistently used within the literature.^{17, 18} Most published studies do not report how participant sex and/or gender information were obtained, nor clearly distinguish between differences related to biological sex or gender. Differences discussed below therefore may represent a combination of differences due to biological sex and gender.

Concussion epidemiology in high performance sports

During the 2019 National Rugby League (NRL) season in Australia, an incidence of 15.4 diagnosed concussions per 1,000hrs match play was reported. During the COVID-19 affected 2020 season, the concussion incidence was 14.9 diagnosed concussions per 1,000 match play hours.¹⁹ In the AFL, the diagnosed concussion incidence for the 2019 and 2020 seasons were 6.5 per 1,000 player hours and 6.8 per 1,000 player hours respectively. In men's rugby-7s match-play, concussion incidence was 3.01 per 1,000 player match hours.²⁰ Injury surveillance data over 6 years from the elite women's rugby union players in England reported 5 concussions per 1,000 hours of match play.²¹ In the Australian Football League Women's Competition, diagnosed concussion incidence for matches from 2017, 2018, 2019, 2020 and 2021 were 15.1, 17.2, 11.5, 4.76, 8.27 per 1,000 player hours, respectively.²² In the National Basketball League (NBA), the annual rate of concussions was approximately 9.5-14.9 recorded between 1999-2018 with the number of games missed increasing from 1.6-3.0 to 3.9-5 following the implementation of the [NBA concussion policy](#).²³

A survey during the 2020 NRL preseason reported that 17% of players did not report a likely concussion to medical staff during the 2018 and 2019 seasons despite 85% of surveyed players receiving concussion education through their club over the previous two seasons.¹⁴ Under-reporting of concussions and failing to seek medical advice range from 17% to 82% across different sports.^{14-16, 24, 25} The fact that large numbers of concussions are going undetected and, therefore, unmanaged is concerning. The reasons players have indicated that they decline to report concussions are primarily due to 'not wanting to be ruled out of the game or training session' (58%) or 'not wanting to let down the coaches or teammates' (23%).¹⁴ These results are consistent with other survey data reviewing the knowledge and attitudes of athletes about concussion.^{16, 24, 26}

In June 2021, Aboriginal and Torres Strait Islander people represented 4% of the total Australian population.²⁷ At the high performance level, First Nation Communities are highly represented per capita. For example 12% of the NRL players, 21% of State of Origin players and 35% of the players in the Australian National team in 2011 identified as Indigenous.²⁸ However, there are no studies comparing rates of concussion between Indigenous and non-Indigenous players. There is need for community codesigned models of research into the epidemiology of concussion in First Nations Communities, as well as an assessment of Aboriginal and Torres Strait Islander peoples' knowledge and attitudes relating to concussion.^{29, 30}

Sex differences in concussion

The popularity of women's contact and collision teams sports is growing. There is a relative lack of research into epidemiology of concussion in female athletes. Evidence supporting sex-based differences in concussion rates is inconsistent.^{9, 31, 32} Females experience worse outcomes from prolonged symptoms compared to males.^{9, 31, 32} Female soccer, basketball, and softball players have an almost two times greater risk of concussion compared to males despite playing under the same rules.³³⁻³⁶ A meta-analysis of concussion incidence in men's and women's rugby 15's revealed contrasting results with an overall incidence of match-play concussion of 4.73 and 0.55 per 1,000 player match hours, respectively.²⁰ In sports like ice hockey and lacrosse, female players are prohibited from intentional body contact (i.e., body checking). Concussion rates are similar between male and female ice hockey players^{37, 38} but female lacrosse players report more concussions.³⁹ Injury surveillance data over 5 years from the National Collegiate Athletic Association (NCAA) Injury Surveillance Program indicates concussion rates per 1,000 athlete-exposures were 6.3 [female] vs 3.4 [male] in soccer/football, 6.0 [female] vs 3.9 [male] in basketball and 3.3 [female] vs 0.9 [male] baseball and softball. In swimming and diving the survey indicated that male rates [0.5] exceeded those of females [0.3].⁴⁰

Many hypotheses exist as to why females experience a higher symptom load and a more prolonged recovery, compared to males. The evidence however is limited. Combinations of intrinsic and extrinsic factors may contribute to these sex-based differences, such as female-specific head impact kinematics, and sex-specific physiological parameters. One hypothesis is a lower biomechanical threshold tolerance for head impacts resulting in more severe symptomology in females.³²

Intrinsic physical factors such as lower neck strength may predispose women to greater head-neck acceleration during impact, increasing the risk of concussion.³² One of the first studies to analyse sex-specific head injury mechanisms demonstrated maximal isometric neck strength of female rugby union players to be 47% lower than males.⁴¹ The same study demonstrated 51% of the head impact events led to whiplash-style head kinematics compared to only one occasion in a male player. In women's collegiate ice hockey, a mean peak linear acceleration of $43g \pm 15g$ was generated by concussive impacts,⁴² significantly lower than concussive impacts in male collegiate players.⁴³ The tolerance for linear head impacts and biomechanical threshold for concussions therefore may be lower in female athletes compared with males.³² The reasons for such differences between sexes are currently not clear. When female soccer players were fitted with a helmetless head impact measurement device, the majority of linear accelerations between 20–40g occurred in practice rather than competition over the course of a season (11% vs 7%, respectively).⁴⁴ Common mechanisms of head injury between sexes also vary. Female soccer players more commonly have concussions through contact with objects (42%) compared to contact with another person (32%).⁴⁵ The opposite trends are seen in male soccer players.

A link is emerging between progesterone levels [i.e., menstrual cycle phase] and risk of concussion and/or outcomes following concussion.^{46–48} For instance, the risk of concussion is high during the late luteal phase of the menstrual cycle [high progesterone concentration] and during the first two days of menstruation.⁴⁷ Further, those who sustained a concussion during the luteal phase of their menstrual cycle, report higher incidences of negative quality-of-life outcomes and neurologic symptoms after one month.⁴⁸ Concussion may also alter hypothalamic-pituitary-ovarian axis function consequently altering menstrual cycle.^{32, 49, 50}

Another hypothesis is that female athletes are more attuned with their bodies than males. Female athletes are more likely to report a concussion than their male counterparts.⁵¹ Therefore, female athletes are more likely to recognise subtle changes/signs and symptoms and more willing to report these changes to team staff than males.⁵²

Research on female concussion is at an early stage and there is significant heterogeneity of data. Extrinsic factors such as rules, use of safety equipment, and variations of training and practices may also contribute to sex differences in concussion. Further knowledge about sex differences in concussion, and both short-term and long-term consequences of concussion is needed. It should be noted that owing to a lack of conclusive evidence, sex is not recognised as a modifier for the management of concussions in the 6th edition of the concussion consensus statement. Rather, sex may be a potential risk factor and/or increase the severity of concussions.⁴

Concussion epidemiology in high-performance para-athletes

The International Paralympic Committee (IPC) define para athletes as a sportsperson with a disability.⁵³ Concussion remains an issue of concern for para-sports due to a lack of research on the validity of concussion recognition tools when applied to a cohort with a wide range of disabilities. Based on experience and available data, the IPC Medical Committee has put together a 'best estimate' risk assessment for concussion which considers impairment, impact speed, collision potential, and use of protective wear. The risk of concussion varies across sports, for instance concussion risk rating for para athletics field (wheelchair) is 1 (low), for para athletics field (amputee) is 2, para athletics track (wheelchair) is 3, Para triathlon (bike, multiple impairments) and football 5-a-side (vision impairment) is 4 and bike and hand cycling, Para alpine downhill (sit-ski, VI and standing) have the highest risk rating of 5.⁵⁴

Table 1: Para sports Summer and Winter with 'best estimate' of concussion risk based upon impairment type, speed, collision potential, protective wear and risk rating: 1 [low] to 5 [high]⁵⁴

Abbreviations: SCI = spinal cord injury, CP = cerebral palsy, VI = visual impairment

Summer Sports	Impairment	Collision Potential	Impact Speed	Head Protection	Risk Rating
Archery	Multiple	Very low	Very low	No	1
Boccia	CP	Very low	Very low	No	1
Cycling Road	Handcycle	Moderate	High	Yes	5
Cycling Road	Trike	Moderate	Moderate-High	Yes	3
Cycling Road	Bike	Moderate	High	Yes	5
Cycling Track	Multiple	Moderate	Moderate	Yes	3
Equestrian	Multiple	Low	Moderate	Yes	2
Football 5-a-side	VI	High	Low	No	4
Football 7-a-side	CP	Moderate	Low-Moderate	No	2
Goalball	VI	Moderate	Moderate	No	3
Judo	VI	Moderate	Moderate	No	2
Para athletics Field	Wheelchair	Low	Very low	No	1
Para athletics Field	Amputee	Low	Moderate	No	2
Para athletics Field	VI	Low	Moderate	No	2
Para athletics Field	CP	Low	Moderate	No	2
Para athletics Track	Wheelchair	Moderate	Moderate	Yes	3
Para athletics Track	Amputee	Low	Moderate	No	1
Para athletics Track	VI	Low	Moderate	No	1
Para athletics Track	CP	Low	Moderate	No	1
Para canoe	Multiple	Low	Low	No	2
Para Powerlifting	Multiple	Very low	Very low	No	1
Para swimming	Multiple	Low	Low	No	2
Para-Triathlon — Bike	Multiple	Moderate	High	Yes	4
Para-Triathlon — Run	Multiple	Low	Low	No	2
Para-Triathlon — Swim	Multiple	Low	Low	No	2
Rowing	Multiple	Very low	Low	No	2
Sailing	Multiple	Moderate	Moderate	No	3
Shooting Para sport	Multiple	Very low	Very low	No	1
Sitting Volleyball	Multiple	Low	Low	No	2
Table Tennis	Multiple	Low	Low	No	1
Wheelchair Basketball	Multiple	Low	Low	No	2
Wheelchair Fencing	T2-T10, ;Below	Low	Low	Yes	2
Wheelchair Rugby	SCI	High	Low	No	3
Wheelchair Tennis	Multiple	Low	Low	No	2

Winter Sports	Impairment	Collision Potential	Impact Speed	Head Protection	Risk Rating
Para alpine Downhill	Sitski	Very high	Very high	Yes	5
Para alpine Downhill	VI	Very high	Very high	Yes	5
Para alpine Downhill	Standing	Very high	Very high	Yes	5
Para alpine Other	Sitski	High	High	Yes	4
Para alpine Other	VI	High	High	Yes	4
Para alpine Other	Standing	High	High	Yes	4
Para alpine Slalom	Sitski	Moderate	High	Yes	3
Para alpine Slalom	VI	Moderate	High	Yes	3
Para alpine Slalom	Standing	Moderate	High	Yes	4
Para cross-country skiing/Biathlon	Sitski	Low	Low	No	2
Para cross-country skiing/Biathlon	VI	Low	Low	No	2
Para cross-country skiing/Biathlon	Standing	Low	Low	No	4
Para ice hockey	Multiple	High	Moderate	Yes	1
Para snowboard	Standing	High	Moderate	Yes	2
Wheelchair Curling	Multiple	Very low	Very low	No	1

During the 2012 Summer Paralympic Games, head and face injuries accounted for 14% of total injuries sustained in 5-a-side football (players have visual impairment) and 7% of total injuries sustained in 7-a-side football (players have cerebral palsy).⁵⁵ Injury surveillance data from the 2016 Summer Paralympic Games also indicates a high frequency of head and face injuries in football 5-a-side (vision impaired football). From total injuries reported at the 2016 Summer Paralympic Games, 8% of injuries were to the neck and 2% were to the head and face. Incidence rates were 0.7 and 0.1 per 1,000 athlete days respectively.⁵⁶ Despite head-to-head contacts being observed on review video footage with players holding their heads with apparent balance issues following the incident, no concussions were reported.⁵⁴

Injury surveillance data from the 2018 Winter Paralympic games showed high frequencies of head and face injuries in Para Alpine Skiing, Para Ice Hockey and Para Snowboarding.⁵⁷ Injuries to the head/face/neck were reported as 20% of all injuries, with 13% of these (i.e., 2% of all injuries) being concussions treated with medical care.

Concussion epidemiology in recreational sports

The United States Centers for Disease Control and Prevention (CDC) estimates that 1.6 to 3.8 million concussions per year result from sports and recreational activities.⁵⁸ Extrapolating from this North American data, it is reasonable to estimate at least 100,000 sport related concussions occur in Australia each year. Many concussions occur in community sport but may go undocumented or fail to come to the attention of healthcare practitioners. Most concussions follow an uneventful course of recovery over days or weeks. There is usually no requirement for medical intervention. However, of the cases that required medical treatment in a hospital, Victorian data demonstrated a significant increase in frequency from 443 per year in 2002–03 to 621 in 2010–11, an increase of 61% over the nine-year period.¹ This was not explained by increased participation, since rates of concussion per 100,000 participants also increased significantly during this time. Hospitalisation rates for concussion across different sports have been examined, and when adjustments were made for participation rates, the sports with the highest concussion rates were determined to be motorsports (181 per 100,000 participants), equestrian (130 per 100,000 participants), Australian football (80 per 100,000 participants), all codes of rugby (50 per 100,000 participants), and roller sports (45 per 100,000 participants).¹

Concussion epidemiology in youth sports

Concussions are common in children and adolescents.^{59–62} By the age of 10 years, one in five children will experience a concussion but only 25% of those concussions result from sport participation.⁶¹ It is difficult to accurately quantify incidence and prevalence in this population due to the lack of data reporting within school sports, lack of awareness of clinical symptoms by parents, teachers and coaches, and youth athletes not reporting head injuries for fear of missing out on playing.^{63, 64} Adolescent males seem to be at a higher risk than females for concussions caused by accidents, not just in sport-related incidents.⁶¹ An Australian national study revealed that 4% of teenage males aged 14–15 years old required medical care for concussions in 2018.⁶⁵ This is higher than females of the same age where only 1% required medical care for a concussion.

Young skulls are large compared to their brains because their brains are not fully developed, therefore easily move within the skull. Young brains have less myelination than adult brains and continue to increase/grow in size throughout adolescence^{66–68} until about 24 years.⁶⁶ Lack of myelination and the potential for brain to move easily within the skull predispose nerve fibres to injury during RHT, making youth more vulnerable to concussion. Weaker neck muscles in youth are proposed as being a confounding factor in impairing the attenuation of forces impacting the head and can increase the risk of concussions [compared to adult populations].⁴⁵

Economic impact of concussion

Aside from health concerns, concussions and traumatic brain injuries represent a significant economic cost to the community. The majority of traumatic brain injuries occur in the 15–64 year age group, representing the group most likely to be engaged in the workforce.⁶⁹ Victorian data estimated the concussion-related hospital admissions cost \$2 million per year,^{1, 70} costing \$1,583 per admission. The Victorian population represents approximately one quarter of the population of Australia. Therefore, extrapolating concussions nationally, the cost to the Australian health system is at least \$50 million annually.⁷⁰ It should be noted that these cost estimates do not account for individuals seeking treatment from primary health care settings such as general practitioners, allied health services, or loss of productive life-years. These figures represent only a portion of the economic costs involved, as they do not reflect costs to the individual, income, and productivity losses, nor that of other regions within Australia.

PATHOPHYSIOLOGY

The pathophysiology of concussion is complex and multifactorial. The physiology underlying concussion symptoms are not yet completely understood. The initial biomechanical impact can initiate a cascade of events resulting in abnormal function at the cellular level. There is also evidence that shearing forces on the brain tissue can alter blood flow to the brain tissue and affect metabolism, affect membrane permeability, release excessive neurotransmitters and contribute to neuroinflammation.⁷¹

Theoretical models are based on animal research and functional neuroimaging studies. Evidence points toward a series of interrelated changes that result in impaired neuronal function^{72, 73}. It is thought that biomechanical forces cause neuronal cell membrane disruption and axonal stretching. The ion channels on the cell membranes become dysregulated and allow an indiscriminate flux of ions. Potassium efflux and calcium influx will result in depolarisation leading to the release of the excitatory neurotransmitter glutamate. In attempting to restore the resting membrane potential, ATP-dependent membrane ion pumps become overactive. This increases glucose demand resulting in a temporary depletion of intracellular energy reserves. There may also be a reduction in cerebral blood flow during this time, further impacting the energy shortage. The intracellular accumulation of calcium due to uncontrolled influx may result in sequestration of calcium within the mitochondria. Mitochondrial calcium overload causes mitochondrial dysfunction, which would further aggravate energy supply issues. These changes are also thought to result in increased free radical production and inflammatory processes, which may be implicated in some of the longer-term symptoms associated with concussion.⁷⁴

During head impact, gelatinous grey matter undergoes significant shear forces from sudden acceleration-deceleration and pulling/pushing of the brain tissue which can cause structural damage. Impact forces can affect the brain in males and females in different ways due to structural differences of brain tissue. For instance, male brains are approximately 10% larger with more global grey matter proportions, whereas, the right hemisphere of the female brain has proportionally more grey matter.⁷⁵ Furthermore, female axons are smaller with fewer microtubules than male axons.⁷⁶ Thus, female axons are at greater risk of failure during trauma under the same applied forces compared to male axons. Further, when exposed to the same mechanical injury, significantly more swelling and greater loss of calcium signalling function can be seen on female axons 24 hours post-injury compared to male axons.⁷⁶

Functional imaging studies have been used to assess physiological alterations underlying concussion and their time course post-injury. Most imaging modality studies suggest long-term alterations beyond the return to normality in clinical and neuropsychological measures. As such, imaging modalities may be useful in research settings to detect changes consistent with concussion and monitor progress beyond recovery of symptoms. However, the current level of evidence in favour of their clinical application is low. Modalities include magnetic resonance spectroscopy (MRS), functional magnetic resonance imaging (fMRI), diffusion tensor imaging, cerebral blood flow measurements, electrophysiology and positron emission tomography (PET) scanning. MRS can detect metabolic changes associated with concussion up to 30 days post-concussion.⁶ fMRI has found differences in functional brain activation patterns from three days up to 23 months post-concussion in concussed athletes compared to controls.⁷⁷ Different diffusion tensor imaging methods have revealed changes in white matter orientation up to six months after concussion.⁷⁸ Alterations in cerebral blood flow resolved 30–40 days post-concussion and electrophysiology did not return to normal levels until 45 days after the injury.^{79, 80} PET scans have revealed changes in cerebral glucose metabolism in some brain regions when comparing military veterans with post-concussive symptoms to controls. This suggests metabolic abnormalities may be implicated in post-concussive symptoms.⁸¹ These neuroimaging modalities suggest the presence of physiological alterations that continue beyond clinical recovery from concussion. However, they are currently limited to characterising the pathophysiology of concussion and are not considered to be tools for clinical assessment. Increased access to high technology modalities, further research and reliability assessment are required before implementation in a clinical setting.

An improved understanding of the pathophysiology of concussion will allow more accurate diagnosis and evidence-based management of the condition. It may provide an enhanced appreciation of the long-term consequences of RHT and concussion and particularly recurrent concussions, to inform risk profiling and mitigation.

RISK REDUCTION/PREVENTION

All sporting organisations have a duty of care to ensure that their sport is conducted in the safest manner possible. Risk of injury, including concussion and RHT, is inherent in many sports, including but not limited to snow sports, cycling, equestrian, contact, collision, and combat sports. It is not possible to remove all risk from sport.

Prevention of RHT in sport is challenging. The main pathways to reduce RHT incidents are via changes to rules/regulations within sport and by modification to training methods to decrease the likelihood of head trauma. Modifying the approach to training, such that there is less emphasis on activities involving potential head trauma, can significantly decrease the overall exposure period during an athlete's career.^{82, 83} In relation to rule modifications, Australian contact sports such as NRL, AFL and Rugby union have modified sporting rules to reduce the risk of concussion in children. The rule modifications are based around reducing the opportunity for high-risk contact. Three main modifications are present and relate to reduced number of players, reduced amount of time spent playing and the reduction of high-risk collisions. Research suggests such strategies may be useful. For example, the removal of body checking in ice hockey⁸² and intentional contact in lacrosse⁸⁴ are associated with a reduction of sport related concussions. Current rule modifications introduced in Australian contact sports work on an incremental process, whereby skills are gradually introduced to children over several years. Research based on tackling has identified that correct education and implementation during training can be associated with a reduced rate of concussion.⁸⁵⁻⁸⁷ Further research is needed to understand the effectiveness of rule modifications that work on an incremental basis, and the long term impacts such changes may have. Furthermore, changes to rules and regulations within a particular sport should be based upon analysis of head trauma risk within the sport and evidence supporting the hypothesis that the rule/regulation changes will decrease risk of head trauma. In December 2023, Rugby Australia announced a two-year trial to lower the legal tackle height from below the shoulders to below the sternum across the community game.⁸⁸ Also in December 2023, the Rugby Football League in Britain announced it will lower the legal tackle height to below the armpit from 2024.⁸⁹ Evidence-based RHT and concussion prevention programs need to be developed, to reduce modifiable risk factors, followed by their implementation in real world settings.

Rule changes also include sports having modifications for games in youth sport, allowing for a minimisation of exposure time and risks that an individual may experience a concussion. Australian contact sports such as [AFL](#), [NRL](#) and [Rugby Union](#) have modifications in place that shorten game times, reduce field size, limit tackle requirements, and change ball size in the under-age categories. This allows for a lower overall exposure to concussion during the formative years of skill development before they are exposed to full competitive contact minutes.

Personal protection equipment (PPE) such as helmets, soft-shell headgear and mouthguards have mixed results in their effectiveness to prevent concussion, in studies to date.⁹⁰⁻⁹³ The use of helmets have been suggested to reduce concussion, with inconclusive evidence. For example research investigating the role of helmet design have identified fit,⁹⁴ type,^{95, 96} and amount of padding⁹⁷ as having a role in concussion detection and symptom development. Likewise, data regarding the effectiveness of soft-shell headgear is also inconclusive; soft-shell headgear has been effective at reducing concussion in adolescent soccer,^{82, 98, 99} with limited effectiveness in other sports such as rugby, and lacrosse.⁸² Finally, the use of protective mouthguards has been effective at reducing the rate of concussion in ice hockey but not in rugby.⁸² Importantly the effectiveness of mouthguards in reducing the rate of concussion does not relate to smart mouthguards which use accelerometer-based technology to measure the impact, intensity and force of head movement in elite athletes. Nonetheless, PPE can play an effective role in reducing head trauma injuries such as lacerations, skull fractures and dental trauma. PPE research continues investigating the use of novel materials but at this stage PPE cannot be recommended for the purpose of preventing concussion.

Several studies have raised the possibility that neck muscle strength is a modifiable factor that influences risk of concussion.^{100,}

¹⁰¹ The evidence regarding the relationship between neck strength and concussion risk is inconsistent but a majority of studies suggest that greater neck strength is protective.¹⁰² Optimising neck strength in those participating in combat, contact and collision sports is a reasonable strategy in concussion prevention.

Beyond prevention, the best way to protect athletes against acute and long-term effects of concussion is to ensure that every concussion is treated seriously and that concussed athletes are removed from the field of play and are not returning to sport prematurely.⁸² A longer symptom free waiting period prior to commencing a return to play protocol leads to a greater reduction in post-RTP injury rates in high-risk sports.¹⁰³ Over the past two decades, regulations regarding stand down times following concussion have evolved in most sports. In the early 2000s, it was common for athletes to return to play on the same day of the concussion. For the past decade, most sports have had post-concussion stand down times of at least 6 days, and longer in children. Recently there has been a move by several sport bodies, including the AIS, to increase mandatory stand down times.

The effects of these evolving regulations on acute and chronic sequelae to concussion, will take years if not decades to be realised. Mandatory stand down times need to strike a balance between being long enough to ensure recovery, but not so long that athletes are disincentivised from reporting concussion. It is important that research continues into the acute and chronic sequelae of RHT and concussion. This CBHPS24 operates on a principle of an 'abundance of caution'. Where there is any suspicion of concussion, an athlete should be removed from the field of play and should not be allowed to return, until cleared to do so by an appropriately trained healthcare practitioner (**'if in doubt, sit them out'**).

RECOGNISE, REMOVE, REFER

When an athlete is suspected of having a concussion, first aid principles apply. A systematic approach to the assessment of airway, breathing, circulation, disability, and exposure applies in all situations. Cervical spine injuries should be suspected if there is any loss of consciousness, neck pain or an injury mechanism that could have led to spinal injury. Manual in-line stabilisation (MILS) should be undertaken until a cervical spine injury is ruled out.

All athletes suspected or confirmed of sustaining a concussion should be immediately removed from the sporting environment and should not be permitted to return to physical activity until they have been assessed by a healthcare practitioner (HCP: an AHPRA registered health care practitioner with appropriate training and experience in concussion assessment and management). Referral to an HCP with appropriate training and experience in concussion assessment and management should occur as a matter of priority. Recognising concussion is critical to correct management and prevention of further injury. Any athlete with a suspected concussion should undergo clinical assessment by an HCP, if possible. Strategies for optimal management of concussion and reduction of recurrent concussion rates include:

- > mandatory removal from sporting environments following actual or suspected concussion
- > adherence to mandatory return to sport protocols
- > requirement to receive clearance to return-to-play from an HCP, and
- > education for key stakeholders such as coaches, parents/guardians/caregivers, teachers and athletes regarding signs and symptoms of concussion.¹⁰⁴

At the high-performance level, some football codes, including the AFL, NRL, and World Rugby/Rugby Australia, have developed criteria for mandatory removal of athletes from sport following head trauma (Note: this policy only applies at the 'elite' level of the sport). These criteria are intended to provide a decision-making support process for doctors to determine the requirement for removal from the sport. The criteria are subdivided into those that require **'immediate removal and no return to sport'** and those indicating **'immediate removal from the sport for further assessment with concussion assessment tools'**.

In recreational sport, there may not be a trained HCP present. In this situation, any suspicion by a match official, coach, athletic trainer, teacher, first aider, or dedicated observer, should result in the permanent removal of the athlete from the field of play. **The Concussion Recognition Tool (CRT6)** is a free resource available to individuals without medical training, to help identify concussions in children and adolescents and adults. Thus, **CRT6** is the most appropriate tool for recreational sport. A concussed athlete is not permitted to return to play that same day and must follow the graded return to sport framework (GRTSF) before being cleared by an HCP. (**'if in doubt, sit them out'**).

In 2018, Rugby Australia introduced the mandatory use of **Blue Cards** in all club, school and domestic representative rugby. When a player exhibits signs and/or symptoms of concussion or suspected concussion, the referee will show the player a **Blue Card**. The Blue Card is a visual cue for team's support staff and triggers an off field medical process to begin. Referees and match officials are one of the 'constants' in the sporting arena. Having these individuals trained to recognise overt signs of potential concussion, to collaborate with team support staff, and empowered to issue a **Blue Card** is highly likely to contribute to the early recognition of concussion, particularly at the community level.

In 2020, the NRL implemented an **On-Field Policy** aimed at community competitions along with a digital Injury Report Form housed on the games registration and competition management platform, **MySideline**. The On-Field Policy dictates what safety personnel and necessary qualifications are required for games to proceed. There are 3 levels of qualification in the game (not accounting for sports physicians at the elite level) being League First Aid, Level 1 Sports Trainer, and Level 2 Sports Trainer, all accreditations include comprehensive training on managing a suspected concussion. At the community level, games for U6-U15 require a minimum of 1 League First Aid per game, for games U16 and above a Level 1 or Level 2 Sports Trainer per team is required or games do not go ahead.

MySideline is used to capture all participant registrations in addition to managing competitions across the country. In the instance of one of the games safety personnel, as determined by the On-Field Policy, suspects a participant may have suffered a concussion, this is included in an Injury Report Form, which in turn automatically renders the participant unavailable for selection for any further games until a medical clearance has been provided to the respective League Administrator.

All sporting codes need to continue exploring strategies for improving systems that enhance safety in relation to concussion and head trauma.

Clinical features of concussion

Concussed athletes may exhibit signs of disorientation, clumsiness, or loss of balance and will often display difficulty concentrating and answering specific questions.⁴ Accurate and timely diagnosis of concussion can be challenging due to the lack of definitive diagnostic markers or investigations and the overlap of symptoms with other neurological, musculoskeletal, and psychological diagnoses. Hence, concussion is a clinical diagnosis — i.e., identifying a concussion is based on the individual's history, symptoms, and signs on physical examination.⁴

A key concept in on field assessment is the rapid screening for a suspected concussion, rather than the definitive diagnosis of head trauma. *The trigger for a concussion evaluation in contact and collision sports across all playing levels is when an athlete experiences a direct or indirect impact to their head and/or trunk associated with visible signs of a concussion and/or the athlete reports symptoms and/or clinical suspicion by medical staff and/or when a possible concussion event is reported by match officials.*¹⁰⁵ **'If in doubt, sit them out'**.

To guide the immediate course of action, mandatory (immediate removal from sport) or discretionary visible signs of concussion have been identified by several sporting codes, for use in high performance athletes.¹⁰⁵ It should be noted that there are minor variations between sporting codes as to what signs of concussion are mandatory or discretionary. Further, there is variation between sporting codes for the course of action following a player displaying mandatory signs of concussion (i.e., immediate removal from the field of play/game, on field assessment and off field management).

Mandatory signs of concussion (immediate removal and no further play)

Athletes displaying any of the following clinical features (i.e., mandatory signs of concussion) should be immediately removed from sport/field of play/game:

- > loss of consciousness
- > lying motionless for >5 s
- > no protective action was taken by the athlete in a fall to the ground, directly observed or on video
- > impact seizure or tonic posturing
- > confusion, disorientation
- > inability to respond appropriately to questions
- > memory impairment/amnesia
- > balance disturbance or motor incoordination (e.g., ataxia)
- > athlete reports significant, new, or progressive concussion symptoms
- > dazed, blank/vacant stare or not their normal selves
- > behaviour change atypical of the athlete.^{10, 105, 106}

When an athlete is removed from the field of play/game owing to mandatory signs of concussion, some sporting codes do not allow that athlete to return to the field of play/game.^{10, 105, 106} In other sporting codes mandatory signs of a concussion is a trigger for mandatory clinical assessment in a quiet and distraction-free environment to determine if the player should return to the field of play/game.^{10, 105, 106}

Discretionary signs of concussion (immediate removal for further assessment)

Athletes displaying any of the following discretionary signs of concussion should be immediately removed from sport/field of play/game for further assessment. Clinical assessment should be conducted in a quiet and distraction-free environment to diagnose or rule out a concussion.^{10, 105}

- > clutching their head*
- > being slow to get up*
- > suspected facial fracture
- > possible balance disturbance or ataxia
- > behaviour change atypical of the athlete †
- > other clinical suspicions.^{10, 105}

* The healthcare practitioner should exercise their medical judgement regarding whether to remove the player for an acute evaluation.

† Some sports consider this a definitive removal criterion.

Red flag signs of concussion (immediate referral to emergency department)

Some features suggest more serious injury and any athlete displaying any of these signs should be immediately referred to the nearest emergency department:

- > neck pain
- > increasing confusion, agitation, or irritability
- > repeated vomiting
- > seizure or convulsion
- > weakness or tingling/burning in the arms or legs
- > deteriorating conscious state
- > severe or increasing headache
- > loss of vision
- > visible deformity of the skull
- > loss of consciousness
- > unusual behavioural change
- > double vision.^{10, 105}

RE-EVALUATE

Symptoms of concussion

Concussive presentations and symptomology may vary significantly and can be complex in nature.¹⁰⁷⁻¹⁰⁹ There are several clinical presentations that can exist following concussion and these can be classified into 5 domains: somatic, musculoskeletal, neurological, fatigue and sleep.¹¹⁰⁻¹¹² Any combination of these domains can form clusters of symptoms. Although evidence in this space is emerging, common symptom clusters have been identified post-concussion. Cognitive and emotional symptoms coexist together most frequently, and fatigue has been associated more commonly with this grouping rather than with physical symptoms.¹¹⁰ Furthermore, there are several systems that can be affected following concussion including cervical/musculoskeletal, vestibular and oculomotor (VOM), autonomic (tolerance to exertion and aerobic exercise), motor and neurological function.^{112, 113} Prolonged recovery and more complex symptomology have been associated with the presence of VOM impairments.^{110, 114, 115} Evidence suggests females tend to have greater symptom severity with prolonged recovery following concussion, compared to their male counterparts.^{114, 116, 117}

The severity of acute and subacute symptoms has been found as the most consistent predictor of slower recovery.¹¹⁸ Subacute symptoms involving headaches or depression are linked to persistent symptoms. The risk of developing persistent symptoms is greater in individuals with a history of mental health problems, attention deficit hyperactivity disorder (ADHD), learning disabilities and migraine.¹¹⁸ Athletes aged between 8 and 14 years of age with diagnosed ADHD or a learning disability can also have worse baseline concussion assessment scores on VOM assessment domains.¹¹⁹

Thorough subjective and objective assessment to identify impairments can help guide rehabilitation improving prognosis, and minimising potential complications such as post-concussive syndrome (PCS).¹¹¹ Tools now exist that can not only help build a clinical diagnosis of concussion but also help differentially diagnose affected domains.

There is currently no conclusive evidence for a modulating effect of sex on outcomes following a concussion. Nonetheless, there is a growing body of evidence pointing to sex differences in symptom patterns, and neurocognitive function.^{17, 33} Although there is a pattern of females reporting greater overall symptoms than males, when examining individual symptoms or symptom clusters, there are mixed findings between the sexes.^{3, 8, 45, 120} Acute symptom scores are higher in females compared to males however no difference exists between the number of symptoms between groups.^{17, 121-127} Male high school athletes reported more cognitive symptom clusters while females reported more somatic and neurobehavioral symptom clusters.¹²¹ Disorientation/confusion and amnesia are the most common primary symptoms reported by males.^{121, 128} Drowsiness and sensitivity to light are the most common primary symptoms reported by females.¹²¹ Another study found that symptoms within the somatic, emotional, and migraine-cognitive-fatigue clusters were more common in females than males.¹¹⁷ For instance, compared to males, "pressure in the head", headache, "feeling slowed down", difficulty concentrating, feeling more emotional, irritability, and sadness are rated higher by females following a concussion.^{124, 129} Females are 2 to 3 times more likely to have migraines, which increases to 3 to 4 times after puberty, than males.¹³⁰ This may explain greater reporting of migraine-cognitive-fatigue symptom cluster in females following a concussion.

Female athletes experience greater neurocognitive impairment following concussion compared to males.^{17, 33} Visual memory composite scores^{125-127, 131} and reaction times are lower in females compared to males following concussion.³² Furthermore, cognitive impairment after a concussion is 1.7 times higher in females compared to male athletes.¹³² Neuropsychological studies generally showed females performing more poorly than males on measures of visual memory following concussion, though this finding was not consistently reported.

It should be noted that the majority of studies discussed above used subjective assessment tools to investigate sex differences.¹⁷ Similarly, owing to lack of conclusive evidence, the 6th edition of the Concussion Consensus Statement did not consider sex difference as a modifier for management of concussions. Rather it was considered a potential risk factor for concussion that may increase the severity of symptoms.⁴ Further research is needed to better understand the interplay between sex hormones, physiological and biological differences and concussion signs and symptoms, risk factors and severity.

Diagnosis of concussion

The diagnosis of concussion should be made by a qualified HCP based on clinical judgement. It should be noted that currently there is no specific diagnostic test that confirms the presence of a concussion. Therefore, in diagnosing concussion, HCP's need to conduct a clinical history and examination across a range of domains including the mechanism of injury, symptoms and signs, cognitive functioning and neurological assessment, including balance testing.^{133, 134} As part of the overall clinical assessment to assess potential concussion, medical professionals can use the [Sport Concussion Assessment Tool 6 \(SCAT6\)](#) for athletes aged 13 years and older.^{135, 136} The diagnostic accuracy of concussion can be significantly increased when the aforementioned clinical features (i.e., mandatory or discretionary signs) are combined with the mechanism of injury.¹³⁷

[SCAT6](#) encompasses an on-field assessment to be used at the time of the concussion, which includes a brief history of the injury, a Glasgow Coma Score and a series of questions known as Maddocks questions. These questions have been validated as an indicator of concussion and are more sensitive in this context than the standard orientation questions.¹³³ The questions assess athlete orientation (in time and place) and they should be preceded by: 'I am going to ask you a few questions, please listen carefully and give your best effort.'

The modified Maddocks questions are:

- > What venue are we at today?
- > Which half is it now?
- > Who scored last in this match?
- > What team did you play last week/game?
- > Did your team win the last game?

The remainder of the SCAT6 is for use off field, in the medical room, or in the consulting room after a referral for a suspected concussion has been made. A possible 22 symptoms of concussion are listed in [SCAT6](#).^{135, 136}

- | | |
|---------------------------|--|
| > Headache | > "Don't feel right" |
| > "Pressure in head" | > Difficulty concentrating |
| > Neck Pain | > Difficulty remembering |
| > Nausea or vomiting | > Fatigue or low energy |
| > Dizziness | > Confusion |
| > Blurred vision | > Drowsiness |
| > Balance problems | > More emotional |
| > Sensitivity to light | > Irritability |
| > Sensitivity to noise | > Sadness |
| > Feeling slowed down | > Nervous or Anxious |
| > Feeling like "in a fog" | > Trouble falling asleep (if applicable) |

[SCAT6](#) is the most widely used concussion assessment tool in the Australian sport sector. It has the greatest diagnostic utility to assess concussion in the first 72 hours/3 days (acute phase).^{136, 138, 139}

Computerised neurocognitive testing can be incorporated in the assessment, but again, it should not be used in isolation.¹⁴⁰ Baseline neurocognitive testing in the pre-season period can be useful for comparison with post-injury scores. Many programs, however, have reference ranges that can be applied in the absence of a baseline test.

It is important to note that [SCAT6](#) was developed in English, which limits its use in culturally and linguistically diverse populations. There is no evidence that [SCAT6](#) is a culturally appropriate tool for Aboriginal or Torres Strait Islander peoples and Australians with culturally and linguistically diverse backgrounds, especially for those individuals whose first language is not English and might have a different second language.^{141, 142}

Concussion assessment in para-athletes

The [SCAT6](#)¹³⁵ and the [CRT6](#)¹⁴³ might be used in the para-athlete population to assess for suspected concussion, but their use may be limited in this population. For instance, one of the assumptions of the [SCAT6](#) tool is that athletes can see, hear, read, and understand information with competence, they have 'normal' baseline speech and language skills, manual dexterity, and range of movement of the cervical spine to enable participation in neurological examination with a comprehensive assessment of mental status, cognitive functioning, gait, and balance.^{4, 136} Yet, para-athletes may experience one or a combination of impairments such as morphological [amputation, dysmelia, congenital deformity], visual impairments, auditory impairments, spinal cord injuries and central neurologic injuries [cerebral palsy, spina bifida, stroke], posing challenges for suspected concussion assessment.¹⁴⁴ For example, male soccer athletes with cerebral palsy report higher total severity scores, more baseline symptoms, worse scores for immediate memory and Balance Error Scoring System (BESS) measures compared to male athletes without a disability on the SCAT3.⁷ Baseline assessments are used to establish a point of reference, and appropriate considerations should be given to baseline cognitive function as well as visual or physical abilities of the para athlete. Periodic baseline pre-participation assessments are recommended in para-athlete populations.¹⁴⁴ It should be noted that baseline scores between different disability groups can significantly vary and in addition para athletes' cognitive function may also vary with the progression of their disability (without a concussive event). Therefore, for effective concussion assessment and management the attending clinician (ideally a team clinician with prior knowledge of the athlete) must have a comprehensive understanding of the preinjury cognitive function and abilities of the para-athlete.¹⁴⁴

It has recently been suggested that the [SCAT6](#) and other guidelines should also include modifications for use with athletes with disability.^{145, 146} Para-athletes are not a homogeneous population and impairments they experience may vary within individual and team para sports. Currently, internationally recognised concussion assessment tools specific to para-athlete populations do not exist and clinicians have highly variable methods for recognition and management of concussions.^{144, 145, 147} The need for adopting and validating internationally recognised tools such as [SCAT6](#) for para athlete groups was raised by the IPC.¹⁴⁵ Therefore, the Concussion in Para Sport (CIPS) Group utilised sections of the [SCAT6](#), '*immediate or on-field assessment*' and '*office or off field assessment*', to develop a traffic light system as a guide to interpreting [SCAT6](#) results for para-athletes with:

- | | |
|--|--------------------------------|
| > [Single] upper limb deficiency | > CP with spastic quadriplegia |
| > [Bilateral] upper limb deficiency | > Dyskinetic CP |
| > [Single] lower limb deficiency | > Ataxic CP |
| > [Bilateral] lower limb deficiency | > Mixed CP |
| > Impaired vision | > Intellectual impairment |
| > Absent vision | > Achondroplasia |
| > Globe absent | > Arthrogryposis |
| > Spinal cord injury (SCI) with quadriplegia | > Polio |
| > SCI with paraplegia | > Muscular dystrophy |
| > Cerebral palsy (CP) with spastic diplegia | > Multiple sclerosis |
| > CP with spastic hemiplegia | > Spina bifida. ¹⁴⁸ |

The traffic light system rates whether additional considerations are required for para-athletes during recognition and assessment of concussion, as well as the return to sport phases. The rating system moves from green = no anticipated considerations, to yellow = potential additional considerations, and red = SCAT6 item not appropriate for para-athletes. Tables of the rating system that are specific to impairments can be found within the supplementary material of the [Concussion in para sport: the first position statement of the CIPS Group](#).

Concussion assessment in the community

At the community level, the athlete should be permanently removed from play if an athlete, coach, first aider/sport trainer, parent/caregiver, match official, teacher or dedicated spotter has any suspicion of a concussion, particularly given HCPs, sideline spotters, and sideline technology may not be available.¹⁰⁵ **'If in doubt, sit them out'**. HCPs often rely on self-reported symptoms to diagnose concussion in individuals who play recreational sports. **CRT6**, a free resource available to individuals without medical training, helps identify concussions in children, adolescents and adults. Thus **CRT6** is the most appropriate tool for recreational sport. The **CRT6** is a simplified summary of the key signs and symptoms and 'red flags' that should raise a concern about a possible concussion. 20 symptoms listed in the **CRT6** are:¹⁴³

- > Headache
- > "Pressure in head"
- > Balance problems
- > Nausea or vomiting
- > Drowsiness
- > Dizziness
- > Blurred vision
- > Sensitivity to light
- > Sensitivity to noise
- > Fatigue or low energy
- > "Don't feel right"
- > More emotional
- > More Irritable
- > Sadness
- > Nervous or anxious
- > Neck Pain
- > Difficulty concentrating
- > Difficulty remembering
- > Feeling slowed down
- > Feeling like "in a fog"

Once a possible concussion is identified, the **CRT6** advises that the individual must be removed from sport immediately and not be allowed to return to activity until they have completed a graded return to sport protocol and are cleared by a healthcare practitioner.^{143, 149, 150} **'If in doubt, sit them out'**. It should be noted that **CRT6** is relatively insensitive to non-specific symptoms.

Health care practitioner concussion evaluation in a controlled office environment

The complex pathophysiological processes of concussion manifest in varying ways across different clinical domains; thus, a multimodal clinical evaluation tool is required to be used in the serial evaluation of an athlete to identify athlete/individual specific/centric intervention and to monitor recovery. Therefore, the **Sport Concussion Office Assessment Tool-6 (SCOAT6)** was developed as a clinical tool to evaluate and manage concussion in the days and weeks following the first 72 hours/3 days (acute period) of injury.^{151, 152} The **SCOAT6** is for use in athletes 13 years and older. The clinical domains included in the **SCOAT6** are:

1. Global symptom scales
2. Cognition
3. Vestibulo-ocular
4. Cervical assessment
5. Neurological examination
6. Autonomic dysfunction
7. Paediatric-specific
8. Balance assessment/postural stability
9. Sleep assessment
10. Depression/anxiety
11. Exercise stress test and
12. Emerging technologies for office assessment

Therefore **SCOAT6** recommends:¹⁵²

- > Word recall and digit backwards tests — The 10-word immediate recall and digit backwards tests should be used; should the athlete find the word recall task too easy (eg, exhibits a ceiling effect), a 15-word list may be used.
- > Measurement of systolic and diastolic blood pressure as well as heart rate in the supine position (after 2 min rest) followed by repeat measures after 1 min standing, accompanied by the recording of any symptoms (eg, lightheaded, dizzy) associated with postural change.
- > Evaluation of the cervical spine.
- > A neurological examination.
- > Timed tandem gait as a single task and a more complex dual task with the addition of a choice among three cognitive tasks.
- > The modified Vestibular-Ocular Motor Screen.
- > Delayed word recall a minimum of 5 min after completion of the verbal list learning and memory tests.
- > Computerised neurocognitive test batteries where available, may also add value.
- > Due to the potential for sleep disturbances associated with concussion, anxiety related to the injury, and the overlap of depression and anxiety symptoms with concussion, inclusion of optional validated sleep screening tool and mental health screening tools was recommended.

The **SCOAT6** is colour coded to:

- > **Blue** — need only be completed at the first consultation
- > **Green** — recommended components
- > **Orange** — optional components.¹⁵²

It should be noted that **SCOAT6** needs to be evaluated and validated.

Concussion assessment in children aged 5–12 years

The **Child Sport Concussion Assessment Tool 6 (Child SCAT6)** was developed as a standardised tool to be used by HCPs to evaluate children suspected of having suffered a concussion.^{135, 153} It is a modified version of the **SCAT6** suitable for children aged 5–12 years. The key differences are that the symptom evaluation is written in language more appropriate for this age group and the severity score is marked out of three rather than six. The **Child SCAT6** also includes a parent's report of symptoms and severity. The **Child SCAT6** lists 21 child reported symptoms and 21 parent reported symptoms. It is recommended that the symptom evaluation be performed with the child in a resting state.¹⁵⁴ It should be noted that due to the questionable reliability and usefulness in young children, the modified Maddocks questions and orientation questions were removed from **Child SCAT6** and the balance examination for children aged 10–12 years includes only the single-leg stance.¹⁵⁴ Further, **Child SCAT6** is most effective in the acute stage following concussion, and up to day 7 post-injury.¹⁵⁵

Table 2: Symptom evaluation for children

Child reported symptoms	Parent reported symptoms
I have headaches	has headaches
I feel dizzy	feels dizzy
I feel like the room is spinning	has a feeling that the room is spinning
I feel like I'm going to faint	feels faint
Things are blurry when I look at them	has blurred vision
I see double	has double vision
I feel sick to my stomach	experiences nausea
My neck hurts	has a sore neck
I get tired a lot	gets tired a lot
I get tired easily	gets tired easily
I have trouble paying attention	has trouble sustaining attention
I get distracted easily	is easily distracted
I have a hard time concentrating	has difficulty concentrating
I have problems remembering what people tell me	has problems remembering what he/she is told
I have problems following directions	has difficulty following directions
I daydream too much	tends to daydream
I get confused	gets confused
I forget things	is forgetful
I have problems finishing things	has difficulty completing tasks
I have trouble figuring things out	has poor problem-solving skills
It's hard for me to learn new things	has problems learning

Individuals without medical training can use the **CRT6** to help identify possible concussions in children and adolescents and adults. The 20 symptoms included in the **CRT6** are listed above.

Clinical features that mandate **'immediate removal and no return to sport'** include:

- > loss of consciousness
- > no protective action in fall to the ground observed directly or on video
- > impact seizure or tonic posturing
- > confusion
- > disorientation
- > memory impairment (e.g., fails Maddocks questions — see below)
- > balance disturbance (e.g., ataxia)
- > athlete reports significant new or progressive concussion symptoms
- > dazed; blank/vacant stare
- > not their normal selves or observed behaviour change.

Where suspicion remains or concussion is confirmed, the athlete *must not* return to sport on the day of injury and *must* seek medical clearance from a qualified practitioner prior to returning to play. **The AIS recommends that the child must be 14 days symptom free [at rest] before return to contact training, and not return to competitive contact sport until a minimum of 21 days from the time of concussion.** The temporary exacerbation of mild symptoms with exercise is acceptable, as long as the symptoms quickly resolve at the completion of exercise, and as long as the exercise-related symptoms have completely resolved before resumption of contact training. **'If in doubt, sit them out'.**

Health care practitioner concussion evaluation in children in a controlled office environment

The **Child SCAT6** is most appropriate for the acute period (72 hours/3 days) while the **Child Sport Concussion Office Assessment Tool 6 (Child SCOAT6)** is a multimodal clinical evaluation tool developed to evaluate and manage concussion in the days and weeks following the first 72 hours/3 days (acute period) of injury.^{156, 157} The **Child SCOAT6** is for children aged 8–12 years. For the tool to be appropriate for children 8–12 years, some components of the adult SCOAT6 required modifications.

The items modified for the **Child SCOAT6** are:

- > **Symptom checklist** — The child and parent symptom reports (the Health and Behaviour Inventory (HBI)) contained in the **Child SCAT6** are modified for the Child SCOAT6 with the addition of 11 new items¹⁵⁸ from the Melbourne Paediatric Concussion Scale. The symptom list from the HBI¹⁵⁹ is scored separately from the additional items to enable comparison to normative data and to previous administrations of the Child SCAT.
- > **Digits backwards** — The first level is a two-digit string, and the final level is a six-digit string.
- > **Days in reverse order** — This is used in children in lieu of months in reverse order. A new timed component is added, such that both time to complete and number of errors are recorded.
- > **Orthostatic test** — The modified methodology described by Pearson *et al* is used,¹⁶⁰ with blood pressure and heart rate recordings after 2 min supine, then 2 min after standing. An optional middle assessment 2 min after sitting is also included.
- > **Dual-task tandem gait** — The cognitive tasks within the SCOAT6 are modified to age-specific tasks.
- > **Visio-vestibular examination** — These tests are used instead of the abbreviated vestibular ocular motor screening. This paediatric-specific methodology has been described by Corwin *et al*.¹⁶¹

Items added to the **Child SCOAT6**:

- > The **PACE Self-Efficacy Questionnaire**¹⁶² is included, and indicates the degree of the child's confidence in their thoughts, feelings and actions affecting recovery
- > The **Symbol Digit Modalities Test**¹⁶³ is included as a measure of psychomotor processing speed.

Several **paediatric mental health** tools are included:

1. **Paediatric Anxiety**^{164, 165}
2. **Paediatric Depressive Symptoms**^{164, 165}
3. **Paediatric Sleep Disturbance**¹⁶⁶
4. **Paediatric Sleep-Related Impairment**¹⁶⁶
5. **The Paediatric Fear Avoidance Behaviour after Traumatic Brain Injury Questionnaire**¹⁶⁶

The **Child SCOAT6** guides clinical management and not every optional component is needed for every consultation, but rather, the healthcare practitioner should use their own clinical judgement on an individual basis to determine which tests and measures are required for each child. The child and parent symptom lists are valuable in assisting in this decision-making process.¹⁵⁶ The tool can also be utilised to assist with the identification of children who may require referral to other specialists for their recovery management.

The Child SCOAT6 is for use in all children aged 8–12 years; however, many items within the tool have not been validated in children with disabilities, or across diverse cultural and language groups.

It should be noted that **Child SCOAT6** needs to be *evaluated and validated*.

Decision trees to support *on field* concussion recognition and management for non-HCP and HCP personnel can be found in Figures 1 and 2 respectively.

Similar decision trees can be found in Figures 3 and 4 to support non-HCP and HCP personnel respectively through the *off field* recognition of concussion.

Figure 1: Non-healthcare practitioner personnel on field concussion recognition decision tree




Figure 2: Healthcare practitioner personnel on field concussion recognition decision tree




Figure 3: Non-healthcare practitioner off field concussion recognition decision tree





Figure 4: Healthcare practitioner off field initial concussion management decision tree



CONCUSSION MANAGEMENT

There is agreement regarding key principles of concussion management policies by a broad array of organisations, including the CISG, American Academy of Neurology (AAN), US CDC, and sporting organisations. The key principles include a brief period (24–48 hours) of relative rest, followed by a supervised and graduated return to cognitive and non-contact physical activity, with no return to contact and collision activities until cleared by a healthcare practitioner.

All individuals dealing with potentially concussed athletes must understand that concussion is an evolving phenomenon. Therefore, signs and symptoms can change or be delayed reflecting the changing underlying physiological injury status of the brain. The diagnosis of a concussion is based on the clinical judgement of a HCP.^{4, 136} In some instances, it will be obvious that there has been a significant injury where the athlete immediately suffers a loss of consciousness, has a seizure or has significant balance difficulties. However, signs and symptoms of concussion can be variable, non-specific, subtle, and may be difficult to detect. Symptoms that are initially subtle can become more significant in the hours and days following the injury and require repeat/serial evaluations. Owing to delays in presentation, it may take up to 48 hours following a head contact to exclude a diagnosis of concussion.¹³⁴ Parents/caregivers, teachers, coaches and attending HCPs need to be alert to behaviour that is unusual or out of character.

The role of physiotherapy in concussion

There is increasing evidence to support the involvement of physiotherapists in the assessment and management of concussion.¹⁰⁷ While ideally, a medical practitioner should be involved early in the clinical management team (CMT), it is often a physiotherapist who will be physically present at a sporting event, recognise suspected concussion and be responsible for removing an athlete from the field of play. Whether as a primary care practitioner in community sport, or as the consistent point of athlete contact in high-performance environments, the physiotherapist's role also places them in a position to observe evolving concussion symptoms.

Most concussion cases are uncomplicated in nature, with 70–80% recovering within expected timeframes and requiring minimal to no input from the CMT.^{108–112} The physiotherapist's role in these cases may involve oversight and guidance through a graded return to sport framework (GRTSF). However, up to 30% of concussion cases are more complex with prolonged symptoms and recovery.^{108, 110, 111} The physiotherapist can play a crucial role in complex case management including further clinical review of cases that do not progress as expected through the GRTS, or who are identified as having vestibular oculomotor (VOM) dysfunction in initial/early assessment post-concussion. Several studies demonstrate appropriate rehabilitation, including cervical and VOM rehabilitation delivers faster symptom resolution, faster return to function and more complete recovery highlighting the value physiotherapists can play in concussion management.^{107, 167–169} The Interdisciplinary care for athletes with concussion (Figure 5) provides a guide that may be helpful for healthcare practitioners involved in the CMT of an athlete post-concussion.

Interdisciplinary care for athletes with concussion


Effective communication and clear CMT role delineation promotes effective and efficient patient-centred care. The structure of the CMT may vary between sport environments, reflecting resource capacity, and the availability and skills of medical practitioners and physiotherapists. Assuming the availability of a medical practitioner, and a physiotherapist with skills in VOM assessment and rehabilitation, Figure 5 illustrates an appropriate system for managing athletes with concussion.

Figure 5: Interdisciplinary care for athletes with concussion

Figure 5, outlines appropriate steps to follow when managing an athlete post-concussion. In an ideal situation, a medical practitioner and physiotherapist with skills in vestibular and oculomotor rehabilitation are part of the clinical management of each concussion case.

Physiotherapists can play a role in post-concussion management through:

- > Supervision and management of a graded return to sport framework (GRTSF)
- > Recognition and identification of vestibular and oculomotor (VOM) impairments
- > Recognition of the deterioration of an athlete's condition
- > Clinical review of complex cases.



*HCP is an AHPRA registered healthcare practitioner with appropriate training and experience in concussion assessment and management

RELATIVE REST

The current principles of concussion management involve a brief period of cognitive and physical rest during the acute period post-injury, followed by a gradual increase in activities of daily living and cognitive activity, prior to a progressive return to sport. Most current evidence supports relative rest during the acute period (24–48 hours post-injury).^{170, 171} In a randomised control trial (45 intervention, 43 control), individuals (age 11 to 22 years) who were prescribed strict rest for 5 days reported more symptoms and recovered more slowly than those who engaged in some physical activity.¹⁷¹ Strict rest beyond the initial period is not recommended.

One of the aims of concussion management is to minimise disruption to learning for children, adolescents, and young adults. There is evidence that abstaining from screen time during the first 48 hours of recovery is associated with a shorter duration of symptoms.¹⁷² As with physical activity, cognitive stimulation such as using screens, reading, undertaking learning activities should be gradually introduced after 48 hours.

REHABILITATION

Throughout the rehabilitation process, consideration of age, para or able-bodied categorisation, sex, cultural and linguistic background, and level of sport is required for each individual athlete to ensure optimal recovery.

As noted above, most concussion cases will follow a relatively uncomplicated rehabilitation and return to life and sport with minimal, if any, intervention being required. However, 20–30% of athletes experience complex symptomology involving multiple domains which can include:

- > greater severity of symptoms
- > longer time to symptom resolution and recovery
- > persistent symptoms.^{107, 111}

All affected domains need to be addressed to achieve complete resolution of symptoms and a successful return to life and sport.¹⁷³

Submaximal aerobic exercises have been shown to lower symptom scores following a concussion.¹⁷⁴ They may not however, have any effect on recovery times.^{107, 173–175} Individuals with a concussion, particularly those with persistent symptoms, may benefit from tailored multimodal interventions for a faster return to sport.¹⁷⁶ Management of rehabilitation is therefore best undertaken by a CMT, where experts in each area assist in the rehabilitation of specific impairments.

It is recommended that a holistic approach, aiming for return to all aspects of life, is used when rehabilitating an athlete post-concussion. This ensures that the five domains i.e., physical, cognitive, emotional, fatigue and sleep are targeted appropriately (Figure 6), as illustrated in Appendix A Physiotherapy guided rehabilitation of concussion.

Vestibular and oculomotor impairments


In addition to the [SCAT6](#), there are evidence-based tools that can assist the CMT (e.g., physiotherapists or medical practitioners) to identify impairments. Tools such as the BESS (the modified-BESS is a component of the [SCAT6](#)),^{116, 117, 177–179} Near Point of Convergence (NPC)^{177, 179, 180} and VOMS^{114, 115, 177, 180–188} can be used to identify VOM impairments in athletes' post-concussion.^{178, 179, 183, 186, 189, 190} A VOMS assessment within 7-days of injury has a moderate prognostic ability to predict a normal recovery in youth athletes.¹⁹¹ However, none of these tools can or should replace clinical judgement when reviewing a concussed athlete. They may however help guide the CMT in the rehabilitation of athletes' post-concussion.

Correct identification of clinical impairments post-concussion can help guide which member of the CMT is best equipped to support rehabilitation.¹¹¹ Clinical guidelines for further review post-concussion (Figure 6), can be a useful tool to guide the clinical review process.

The combination of validated symptom questionnaires and clinical tools assessing physical impairments (Figure 6) can guide management and referral onto appropriate practitioners within the CMT.^{110, 111, 184} Various symptom identification questionnaires have been developed for adult populations including the Post-concussive Symptom Questionnaire (PCSQ) and its shorter form the PCS-19.¹⁹² The Melbourne Paediatric Concussion Scale (MPCS) is the first questionnaire developed and validated specifically for paediatric populations. The MPCS combines the 21-item Post Concussion Symptom Inventory (PCSI) with 10 expert agreed-upon clinical questions.¹¹¹

Figure 6: Clinical guidelines for further review post-concussion

Initiation of a graded return to sport (GRTS) is recommended following a concussion. Prolonged or severe exacerbation of symptoms that do not return to baseline for the concussed individual can result in failure to progress through the GRTS. In these situations or if vestibular and oculomotor (VOM) impairments are suspected clinical review with an appropriately trained physiotherapist as part of the clinical management team (CMT) is recommended. Below is a guide for physiotherapists and medical practitioners involving specific assessment of cervical, VOM and autonomic systems. Physical assessment and symptom identification questionnaires can be helpful to identify affected symptom domains, guide rehabilitation and ensure complete recovery post-concussion.



RETURN TO SPORT FOR COMMUNITY AND YOUTH SETTINGS

A collaborative multidisciplinary approach to concussion management with shared decision making is encouraged but is not always available outside of an elite, high-performance setting. Children and adolescents have growing brains and are more likely to have a prolonged recovery after concussion.¹⁵⁴ All athletes under the age of 19 should follow a more conservative return to sport/learn protocol. Community and youth environments often have inadequate availability of appropriately trained HCP support to closely supervise individuals through the recovery and return to sport/learn process. This may include schools, local sporting clubs and other non-sport related settings.

As knowledge of concussion management improves, more sporting organisations are opting for longer stand down periods, before return to contact or collision activities is permitted. The updated graded return to sport framework (GRTSF) for community and youth (Figure 7) assists athletes/coach/parents/teachers with concussion management through the recovery process and time frames for a safe return to sport/learn.

The AIS return to sport protocol for community and youth sport includes;

- > Introduction of light exercise after an initial 24–48 hours of relative rest
- > Several checkpoints to be cleared prior to progression
- > At least 14 days symptom free (at rest) before return to contact/collision training. The temporary exacerbation of mild symptoms with exercise is acceptable, as long as the symptoms quickly resolve at the completion of exercise, and as long as the exercise-related symptoms have completely resolved before resumption of contact training.
- > A minimum period of 21 days until the resumption of competitive contact/collision sport
- > Consideration of all domains throughout the recovery process.

Reintroduction of daily activities is appropriate if the activities do not severely exacerbate symptoms following the initial 24–48hour period of rest.^{170, 171} Early resumption of activities of daily living is associated with improved symptom resolution and shorter recovery time.^{107, 171}


After the initial period of relative rest, graded return to school and/or work is advised. Concussion is an evolving injury and symptoms can change over time in one or more domains. All affected domains may not be evident during the early stages of the graded return. Care should be taken when returning to activities that involve multiple domains, such as school or work, with dosage and environment considered.^{107, 112}

Recent studies encourage initiation of low-intensity exercise 24–48 hours post-concussion irrespective of the presence of low-level symptoms.^{4, 177, 193} Mild and brief exacerbation of symptoms may occur during progression through the GRTSF. This is acceptable as long as the exacerbations are temporary, that is, the symptoms return to baseline before the next exercise session.¹⁹⁴ Mild is defined as an increase of no more than 2 points on a 10 point scale and brief is no more than 1 hour duration of exacerbation from pre-exercise symptoms.¹⁰ If there is moderate or severe exacerbation of symptoms or symptoms persist until the next scheduled bout of activity (considered prolonged symptoms) then a review with a HCP is recommended.¹⁹⁴ Earlier clinical review may be warranted if VOM dysfunction is suspected. Reviewing the athlete allows symptom domains that may be affected to be identified and appropriate intervention to be implemented.

Children and adolescents take longer to recover from concussion than adults.¹⁵⁴ A more conservative approach should be taken with those aged 18 years or younger. **The GRTSF requires those under 19 years of age and those without a dedicated HCP to guide recovery, to be symptom free for 14 days [at rest] before return to contact training, and not return to competitive contact sport until a minimum of 21 days from the time of concussion.** To be clear, that is not 14 days from the time of concussion. It is 14 days from when the athlete becomes symptom-free. The day of the concussive incident is deemed day 0 of the GRTSF. This recommendation allows for the individual case variability in symptom duration. It ensures that the most vulnerable individuals have demonstrated a clear capacity to perform all normal activities of daily living, including non-contact exercise, without symptoms, before they return to the field of play.

Figure 7: Graded return to sport framework for community and youth

Each stage, highlighted in orange or green below, should be at least 24 hours and symptoms should return to baseline prior to commencing the next activity or stage.



Some high-performance athletes may have access to appropriately trained Healthcare Practitioners experienced in multi system concussion rehabilitation. These athletes may be cleared earlier if their sports concussion protocol allows. Refer to the graded return to sport framework for advanced care settings. Note, athletes aged under 19 years should NOT have access to earlier clearance available in advanced care settings.

Concussion Officer for Community/School Environments

All community members should be aware of the concussion management protocols and pathways relevant to their sport and community, including their role in the identification and management of concussion. It is recommended that clubs and schools introduce a 'concussion officer' to oversee the management of concussion. A 'concussion officer' is a single point of contact and manages the coordination of matters related to concussion. A 'concussion officer' is not a concussion expert and is not expected to diagnose concussion. Analogous to the role of a 'fire warden', the 'concussion officer' ensures that anyone diagnosed with concussion follows the organisation's agreed concussion protocol. The designated person can be any member of the affiliated community. Their job is to be the recipient of information in relation to concussion and to ensure that the concussion protocol is enacted. It is recommended that a 'whole of community' concussion policy or protocol is adopted. This needs to be effectively socialised and communicated to all stakeholders, to ensure it is adapted successfully.

Figure 8: Systems for managing the concussed student in the school environment





Figure 9: Systems for managing the concussed student in the school environment

Appoint a 'Concussion Officer'

- > Similar concept to a fire warden
- > Role is not to diagnose or provide medical advice
- > Once concussion is suspected / diagnosed, relevant stakeholders are informed
- > Ensures 'whole of school' concussion plan is enacted.



ADVANCED CARE RETURN TO SPORT FRAMEWORK

Advanced care settings (ACS) are defined by the level of HCP support available to athletes. Athletes must have daily access to HCPs that have specific training and experience in the management of concussion and head trauma. The advanced care return to play framework applies to athletes 19 years and over who are in an 'elite' athlete environment. This framework is never available to athletes under 19 years of age. A more conservative approach should always be taken for youth athletes. **'If in doubt, sit them out'**.

To be eligible for an athlete to progress through an advanced care protocol, the ACS should have the ability to provide:

- > Baseline **SCAT6** and/or neurocognitive testing, conducted prior to the injury.
- > Sideline care for training and matches.
- > Video recording of matches for review (and contact training if possible).
- > Access to a CMT to supervise the return to play and instigate any treatment or investigation required should the RTS progression not be straightforward.
- > A formal and documented concussion education program for coaches and players in the club or team involved.

While the general principals of the community based GRTSF described above apply, the graded return to sport framework for advanced care settings (Figure 10) supports eligible athletes. Athletes who have appropriate CMT support, may not need to spend as much time in each recovery stage. Athletes may progress to checkpoint 1 from day 6 and checkpoints 2 and 3 from day 10. Importantly, this framework is not in place to say all athletes will return on a specific day post-concussion, rather the progression through the stages is based on the individual experience and should always be guided by the CMT and the individual athlete.


Protocol for those with multiple suspected concussions

An athlete with a history of multiple concussions is at risk of experiencing prolonged symptoms before return to sport.¹⁹⁵ Those who suffer from multiple concussions within a short period of time should be managed more conservatively and be assessed by a CMT with specific training and expertise in concussion. Multiple concussions can be two concussions within a 3-month period, or three concussions in a 12-month period. If this occurs, the individual should follow a more conservative return to sport protocol. There is no evidence regarding specific time frames for return to sport following multiple concussions. The timeframes will be influenced by factors such as the severity of the most recent injury, the number of previous concussions and the general medical history of the athlete. A recommended starting point for return to sport after second concussion within three months, would be 28 days symptom-free before return to contact training and a minimum of six weeks from the time of the most recent concussion until return to competitive contact. Being part of an "advanced care setting" does not exclude the individual from following a longer recovery period.

In situations where three concussions have occurred within a 12-month period, consideration needs to be given to missing a season of contact / collision sport. These decisions can have serious ramifications for the athlete's livelihood and mental wellbeing. The decision about standdown periods needs to be made after thoughtful deliberations between the athlete, their support persons and the CMT. The discussion should be transparent to highlight the scientific facts and uncertainties of their condition, weighed against the potential risks of returning to their previous sport and/or alternative activities with less chance of concussion.¹⁹⁶

Figure 10: Graded return to sport framework for advanced care settings

Each stage, highlighted in orange or green below, should be at least 24 hours and symptoms should return to baseline prior to commencing the next activity or stage.



EMERGING TOOLS FOR CONCUSSION DIAGNOSIS

Emerging tools

Detecting a concussion in routine brain imaging is difficult as it is predominantly considered a functional neurological disturbance rather than a structural injury. Currently, there is a growing interest in Point of Care (PoC) devices that use biomarkers to provide an objective assessment tool to assist with concussion diagnosis and clinical decision-making. For instance, in a prospective observational study of 1,028 male professional players, salivary small non-coding RNAs (sncRNAs) were identified as unique signatures of concussion.^{197, 198} There is also research underway to explore the potential clinical utility of blood biomarkers as an objective PoC to diagnose concussion.^{128, 199–202}

Although, imaging modalities may be useful in research settings to detect changes consistent with concussion, current evidence does not support the clinical use of these modalities to diagnose or manage concussion.

Given the rapid technical advancements, prudent use of technological advances may improve the accuracy of concussion assessment. However, caution needs to be exercised when using such tools and validation is required before their global adaptation. At present, the evidence base is insufficient to recommend the routine use of any medical imaging or biomarker tests in the diagnosis and management of concussion.

Vestibular and oculomotor assessment

Technologies such as eye tracking devices, neurocognitive tablet applications (i.e., Apps), and other sensory organisation testing devices may have a role in VOM assessment. Access to such tools may vary however, and there is limited evidence for clinical utility in athletic populations. Furthermore, due to the limited accessibility to specialist equipment, many studies are conducted with small sample sizes which can limit applicability of findings in the general population. Ongoing research in this area is required to achieve greater validity and reliability of these tools to be used in the diagnosis and management of concussion.

CONCUSSION AND LONG-TERM BRAIN HEALTH

There is concern about potential long-term consequences of concussion or RHT resulting from ongoing participation in contact, collision, and combat sports.^{203–205} There is an association between a history of exposure to RHT and cognitive deficits later in life.²⁰⁶ There are a growing number of cases of individuals who are posthumously diagnosed with *Chronic Traumatic Encephalopathy Neuropathological Change* [CTE-NC], following a sporting career involving RHT.²⁰⁷ *"A challenge of evaluating the long-term consequences of repetitive head impacts is that the outcomes are chronic, but the exposures are acute and, in this setting, remote. Each impact is of short duration, can be ambiguous, and rarely quantified"*²⁰⁶ It should also be noted that in the 80's, 90's and early 2000s, it was common for athletes to return to play on the same day of the concussion, or to not be removed from play following a concussion. For the past decade, regulations regarding stand down times following concussion have evolved in most sports.

CTE-NC is defined by the accumulation of hyperphosphorylated tau (p-tau) in neuronal cell bodies and processes, located around small blood vessels, at the depths of cortical sulci.²⁰⁸ As with many neurodegenerative conditions, there is an imperfect correspondence between neuropathology and clinical phenotype.²⁰⁷ The term CTE-NC has more recently been adopted to distinguish the neuropathology from associated clinical symptoms (referred to as Traumatic Encephalopathy Syndrome, TES). CTE-NC can only be confirmed post-mortem, based on histopathological examination of brain tissue at autopsy.^{206, 209} In 2021 researchers published the following;

- > a revised consensus criteria for classifying the neuropathology of CTE-NC²⁰⁸
- > the first-ever consensus criteria for diagnosing TES, the clinical condition believed to be caused by repetitive mild neurotrauma.²¹⁰

While CTE-NC is a neurodegenerative pathology associated with a history of RHT, there currently is a lack of high quality evidence indicating the **degree of association** between RHT, concussion in contact and collision sport with CTE-NC, a condition with broad clinical diagnostic criteria (i.e., TES).²⁰³ Some studies suggest a 'dose effect' of repeated concussions increasing the risk of CTE-NC.^{211, 212} The evidence linking RHT and concussion with CTE-NC consists of case reports, case series, and retrospective and post-mortem analyses. Due to the study design, and the reliance on retired athletes volunteering for an autopsy diagnosis, there is significant selection bias in many of the reports.^{204, 213} The research data on CTE-NC is usually obtained from sport brain bank studies. Those who donate their brain for these studies almost universally have pre-existing clinical symptoms of degenerative brain disease. The brain donations are made in good faith but the skewed representation makes it difficult to apply the findings to the general population.²⁰⁴ The clinical data obtained from sport brain bank studies has also largely relied on retrospective interviews with athletes and athlete-relatives for information such as playing time, RHT and exposure, symptom patterns, mental health issues and substance abuse. Recall bias is highly likely to affect the reliability of such information, as is the case with research into other forms of degenerative brain disease.²⁰³ Retrospective clinical analysis is insufficient for creating robust clinical diagnostic criteria for CTE-NC in living patients.²¹³ CTE-NC is not an inevitable consequence of RHT.²¹⁴ Studies to date have not adequately controlled for the potential contribution of confounding variables such as alcohol abuse, drug abuse, personality factors, genetic predisposition, education exposure, family history of mental health and neurological problems.²⁰³

A recent study used the Bradford Hill criteria to review the evidence for the link between RHT, concussion and CTE-NC.²¹⁵ The authors concluded that **"we found convincing evidence of a causal relationship between [RHT] and CTE, as well as an absence of evidence-based alternative explanations."**²¹⁵ It should be noted however that Sir Austin Bradford Hill CBE FRS himself emphasised that this method of analysis does not remove the requirement to differentiate between correlation and causation.^{216, 217}

A large-scale study of 636 cases of community-based cohort of ageing and neurodegeneration from the Sydney Brain Bank collection, CTE-NC was identified in five cases (prevalence 0.8%). Three of the five cases with CTE-NC had a history of traumatic brain injury and two cases had no known history of neurotrauma (including repetitive head impacts from sports).²¹⁸ Low prevalence of CTE-NC is further corroborated from similar sample of 532 cases from the United States and 323 cases from Europe, where CTE-NC was 0.6%²¹⁹ and 0% respectively.²²⁰ This low prevalence of CTE-NC in the general community highlights the need for further well-structured longitudinal studies exploring the strength of the link between RHT, concussion, and CTE-NC.

The fundamental concern regarding CTE-NC is that individuals who have participated in contact and collision sports have experienced damage to the microstructure of their brains, through repetitive neurotrauma,²²¹ which places them at future risk for accelerated brain aging, and neurodegenerative diseases — including mild cognitive impairment, Parkinson's disease, amyotrophic lateral sclerosis, CTE-NC, and Alzheimer's disease.²²² With respect to cognitive changes, a combination of factors such as frequency of RHT and/or concussion, severity, timeframe, age, and sex might predict cognitive changes

better than frequency of RHT/concussion alone.^{148, 223} The degree of cognitive reserve or brain reserve has been shown to protect against brain injury or disease.^{224–226} For example, when an individual is completing a task with increased difficulty, cognitive reserve mediates the establishment of efficient neural networks (i.e., recruiting new and/or less injured networks) to complete the task.^{224, 226}

Recent descriptions of the clinical features of TES has proposed "neurobehavioral dysregulation" (e.g., 'poor regulation or control of emotions and/or behavior, including explosiveness, impulsivity, rage, violent outbursts, having a short fuse or mood swings')²¹⁰ Cognitive impairment, typically affecting memory and executive function has been proposed to emerge later,^{210, 227, 228} and has been described as being 'progressive'.^{210, 227, 228} Studies reporting evidence of an association between elite male sport participation and long-term cognitive, psychiatric, and neurobehavioural problems^{229, 230} invite the hypothesis that such neuropsychiatric features might reflect the early manifestations of CTE-NC.

A number of studies indicate that elite athletes enjoy greater longevity.^{37, 231, 232} The death rates of 9,932 current and retired elite male Australian football players was 21 % lower than the death rates of aged-matched males in the general population.²³³ Similarly, a large-scale retrospective epidemiological study of 3,439 retired male National Football League players (played from 1959 to 1988) reported lower risk for suicide compared to men in the general population.²³⁴ In contrast, another study reported a disproportionate number of suicides over the period from 2009 among current and former professional football players.²³⁵ The causes of suicide are complex. The evidence to support the causal relationship between contact and collision sports, RHT and/or concussion, depression and suicide and CTE-NC is inconclusive as no data from peer-reviewed cross-sectional, epidemiological or prospective studies are available.^{236, 237}

Several studies are being conducted around the world investigating the brain health of former collision sports athletes. In Australia, the *Former Elite Level Athlete Brain Health Research Program*, is a prospective, longitudinal clinicopathological study, that was established in 2012 to investigate variables that affect brain health and aging to evaluate the extent to which modifiable risk factors may contribute to the risk of poor health outcomes. This research program includes elite level former rugby league and rugby union players and comparison groups that consist of age and education-matched healthy community-based control participants without a history of neurotrauma or contact sport participation and former Australian able-bodied Olympians from non-contact, collision, or combat sports (for example, swimmers, rowers, and track and field athletes). The evaluation includes several self-report health questionnaires, a clinical interview, an in-person evaluation (including cognitive testing and brain MRI), and a brain donor program. To date, preliminary data published from this research program has revealed no significant differences between groups (former players vs control groups) on measures of depression, anxiety, or cognitive functioning.²³⁸ Within the player group, smaller subcortical volumes were significantly associated with greater alcohol consumption. Alcohol Use Disorders Identification Test (AUDIT) score was higher in former rugby players compared to controls.^{239, 240} Extending earlier findings from this cohort, a group of 141 former rugby league players (median age = 52.6 years [13.8], range = 30–89 years) who played professional sports for a median of 8 years playing [interquartile range (IQR) 3.5–11] reported a median of 15 lifetime concussions [IQR 6–30].²⁴¹ 29% of former rugby players included in the study reported at least mild current depression. Depression, Anxiety, and Stress Scale (DASS) scores were not significantly correlated with lifetime concussions. Multiple regression analysis revealed that 35% of the variance in DASS depression was accounted for by 'age, concussion history, years played professionally, the CD-RISC, Brief Pain Inventory-pain interference score, and Epworth Sleepiness Scale score.²⁴¹ No relationship was found between objective cognitive test performance and perceived cognitive decline in 135 retired players (mean age 53.1 years [13.9], range 30–89 years). Depressive symptoms were the strongest predictor of perceived cognitive decline.²⁴² No significant difference in the size of the brain ventricles was observed between 41 retired players and 41 controls. Further, there were no significant differences between those with and without an abnormal cavum septum pellucidum and age of first exposure to rugby league, the number of lifetime concussions, years of exposure to repetitive neurotrauma, perceived cognitive decline, cognitive decline on any neuropsychological test, depression, or impulsivity.²⁴³

These findings should be interpreted with consideration for the representativeness of the sample, the sample size, the generational/era differences of the study participants (i.e., the game play of the sport, the physical demands of the sport, the professionalism and training commitments, and injury management differ greatly across generations), self-selection of 'super controls,' that may not reflect the population base rates of condition/disease/illness, and the role of comorbidities and/or confounding variables (e.g. acting as moderators, mediators of various health outcomes).

Long-term brain health in female athletes

Many sports have recently established professional female leagues, providing female athletes with a pathway into elite sports, including contact and collision sports. With this growth in female sports, there is a need to evaluate the long-term brain health in female contact and collision sport athletes. Currently, there is a lack of research on how the female brain responds to concussion and whether the response differs from that observed in males. There is also a lack of research investigating the long-term brain health of female athletes. In the general population, compared to men of the same age the incidence of dementia (85 years) and Alzheimer's disease (80 years) is greater in women.²⁴⁴ Female axons are at greater risk of axonal damage during trauma which may trigger Amyloid- β peptide production and the accumulation of amyloid pathology, a core pathological feature of Alzheimer's disease.²⁴⁵ Although the reasons are unclear, the prevalence of autoimmune disorders amongst women is greater, occurring at a rate of 2 to 1.²⁴⁶ The Amyloid- β peptide plaques that cause Alzheimer's disease, are an innate immune protein activated as part of the brain's immune system to fight infections.²⁴⁷ The female immune system response is stronger compared to males²⁴⁸ and females may therefore have more Amyloid- β peptides, which may increase their risk of Alzheimer's disease. The role of biological differences in women, and any associated effect on predisposition to TES, CTE-NC or other neurodegenerative diseases, remains unclear.

Long-term brain health in para-athletes/athletes with disabilities

It is important to consider the unique environments that para-athletes navigate. To date there are no reports in the medical or scientific literature on the long-term brain health of para-athletes exposed to RHT and/or concussion. Assessment of brain health changes in athletes with a disability presents a unique challenge because of the various para-classifications (e.g., athletes with neurological conditions, learning disorders, and/or intellectual disability).

HOLISTIC BRAIN HEALTH AND WELLBEING IN CURRENT AND FORMER ATHLETES

The long-term brain health and wellbeing of athletes both during, and in the years following retirement are of concern. It is important that all sporting organisations recognise and understand all the factors that potentially contribute to long-term brain health. The Lancet Commission lists contributing factors to an individual's risk of developing dementia, with TBI increasing the risk by 3%.²⁴⁹ Other potentially modifiable factors that need to be considered by athletes and sporting organisations include lesser education (7%), alcohol (1%), smoking (5%), depression (4%), physical inactivity (2%) and social isolation (4%).²⁴⁹

Cognitive reserve refers to an individual's adaptability of functional brain processes (i.e. flexibility, efficiency, capacity, compensation), with individual differences in cognitive reserve allowing some individuals to cope better with age and disease related brain changes.²⁵⁰ A range of studies have demonstrated that cognitive reserve is very important in determining whether a range of histopathological changes of degenerative brain diseases cause symptoms during life. *"As with many neurodegenerative conditions, there is an imperfect correspondence between neuropathology and clinical phenotype"*.²⁵¹ An analysis of clinical records and the pathology of brain donations from 180 patients from two 'Brains for Dementia Research' cohorts found that more than a third of cases clinical diagnosis was different from final neuropathological diagnosis.²⁵² A longitudinal study collecting clinical and neuropathological data on 498 Catholics sisters found that 12% of participants who displayed no clinical symptoms were found to have Braak stages V–VI pathology,²⁵³ demonstrating that pathology can be present without the clinical symptom presentation during life.

There is abundant evidence that cognitive reserve is modifiable and can be optimised with appropriate strategies. The level of education an individual has is linked to a lower risk of Alzheimer's disease.²⁵⁴ Clinical-pathologic studies have shown that lower education is associated with cognitive decline or dementia regardless of Alzheimer's disease pathology.^{255–257} Furthermore, participation in physical, cognitive, and social or leisure activities, has been linked to a reduction of risk for dementia.^{258–260} Involvement in these activities assists individuals through cardiovascular benefits of physical activity and through cognitive stimulation which builds cognitive reserve. Sport organisations are encouraged to adopt a holistic approach to the long-term brain health of current and former athletes. Sporting organisations should provide education for athletes, about how to optimise cognitive reserve and diminish their chances of poor brain health in later life. Figure 11 shows factors linked with cognitive reserve, providing an overview of holistic strategies sporting organisations could target during educational sessions.

More research is required that considers holistic cognitive health in diverse samples of elite athletes and sports while controlling for the impact of exposure to concussions throughout a career.

Figure 11: Holistic brain health



QUESTIONS TO BE ANSWERED IN IMPROVING UNDERSTANDING OF CONCUSSION AND LONG-TERM BRAIN HEALTH

Knowledge regarding the effects of RHT, concussion and long-term brain health continues to evolve. Many questions, however, remain unanswered. Well-structured scientific investigations are needed to address knowledge gaps. The AIS encourages co-design research models incorporating athlete voices and voices of underrepresented communities. Future research should be targeted to answer the following questions;

- > What is the prevalence of CTE-NC in female, male, and para-sport athletes?
- > What is the strength of the association between RHT, concussion, and development of CTE-NC?
- > What is the strength of association between histopathological changes of CTE-NC and the clinical syndrome of TES?
- > Which athletes are susceptible to development of CTE-NC, and why?
- > What role do modifying factors play in susceptibility to the development of CTE-NC?
- > What is the natural history of CTE-NC? Is it an inexorably progressive disease, similar to neurodegenerative diseases such as Alzheimer's disease?
- > Are female athletes more susceptible to CTE-NC than males, for a set dose of RHT exposure?
- > Are para-athletes more susceptible to CTE-NC than able-bodied athletes, for a set dose of RHT exposure?
- > Are specific cultural cohorts more susceptible to CTE-NC than athletes from Anglo-Saxon background, for a set dose of RHT exposure?
- > What are the sex-based differences in risks and clinical picture of RHT and concussion in sport?
- > What is the prevalence of RHT and concussion in First Nations Communities and culturally and linguistically diverse populations?
- > What changes can be made to better capture nationwide data on RHT and concussion in sport?
- > What sport-specific interventions are efficacious in preventing RHT and concussion?
- > What is the impact of RHT and concussion on developing brains of youth athletes and the long-term health consequences?
- > What is the long term mental and physical health in those exposed to RHT and concussion?
- > What is the effect of more conservative return to sport protocols on the acute and long term sequelae of concussion?
- > What are the most effective therapeutic interventions for recovery from episodes of concussion?
- > What rule/regulation modifications could be effective in reducing incidence of RHT in individual sports?

These unanswered questions present an enormous challenge to the medical and scientific community. Prospective, longitudinal clinicopathological studies can help identify possible early clinical features, progression [if CTE-NC is a progressive disorder], and potentially help with interventions.

EDUCATION

General awareness and knowledge about concussion, although improved over recent years with the availability of guidelines and educational materials, remains less than optimal.²⁶¹⁻²⁶³ To effectively improve awareness and understanding in the community, education must be targeted at groups that are at risk of concussion.^{261, 264} The [Concussion in Sport Australia](#) webpage has relevant information for each key stakeholder group.

Athletes — need to have a good understanding of concussion to appreciate the importance of reporting symptoms and complying with rest and return to sport advice.²⁶⁵

Parents/guardians/caregivers, coaches, and teachers — must be able to recognise symptoms and signs of concussion to improve detection at the community level, where there is limited medical supervision present.^{263, 266}

Sporting and medical organisations — need to continue to develop specific recommendations around concussion to inform their members/participants. However, complexity of the return to sport protocol has been highlighted as a potential barrier for community sport.²⁶⁷

One of the key aims of this CBHPS24 is to take the latest information from medical and scientific journals, synthesise the information into an easily digestible format and make the information available to all Australians involved in sport or who have an interest in RHT and/or concussion.

Education Modules

General Population: [Connectivity](#) is a not-for-profit organisation working to help improve the lives of people with traumatic brain injuries. Connectivity has a 15-minute interactive online [Sport-Related Concussion Short Course](#) which aims to educate athletes, coaches, umpires, staff, volunteers, parents/guardians/caregivers and other interested people to recognise and manage a sport-related concussion injury.

Healthcare Practitioners: The AIS, in partnership with Sports Medicine Australia, Australian Physiotherapy Association and Australasian College of Sport and Exercise Physicians has a freely available education module targeted for healthcare practitioners [HERE](#). Healthcare practitioners is defined as AHPRA registered individuals with appropriate training and experience in concussion assessment and management.

KEY POINTS FOR ATHLETES, COACHES, PARENTS, TEACHERS AND SUPPORT STAFF

- > Concussion is a type of brain injury that occurs from a knock to the head or body.
- > Recognising concussion is critical to ensure appropriate management and prevention of further injury.
- > The [Concussion Recognition Tool 6 \(CRT6\)](#) is recommended to help recognise the signs and symptoms of concussion. [CRT6](#) can be downloaded [here](#)
- > First aid principles apply in the management of an athlete with suspected concussion. This includes observing first aid principles for protection of the cervical spine.
- > Any athlete suspected of having concussion should be removed from sport and not allowed to return to sport until cleared by a health care practitioner who has appropriate training and experience in concussion assessment and management.
- > Features that suggest more serious injury and prompt immediate emergency department referral include neck pain, increased confusion, agitation or irritability, repeated vomiting, seizure, weakness or tingling/burning in the arms or legs, reduced level of consciousness, severe or increasing headache, or unusual behaviour.
- > There is no single test that can determine whether someone has sustained a concussion.
- > When assessing a patient with suspected concussion, a healthcare practitioner will ask about details of the event, past medical history, then assess the patient. This can include asking about symptoms, signs, testing memory function and concentration, balance, and neurological function.
- > Once diagnosis of concussion has been confirmed, initial treatment for concussion is relative rest. After 24–48 hours of rest, light intensity physical activity is indicated, that does not cause a significant and sustained deterioration in symptoms.
- > An active recovery following the Graded Return to Sport Framework (GRTSF), for both the community/youth and advanced care setting, should proceed after the initial period of relative rest. Each stage outlined requires a minimum of 24 hours and progression should not be made unless symptoms have returned to baseline for the concussed athlete.
- > Any athlete under the age of 19 should complete the more conservative GRTSF for community and youth. ***This requires those under 19 years of age and those without a dedicated HCP to guide recovery, to be symptom free for 14 days (at rest) before return to contact training, and not return to competitive contact sport until a minimum of 21 days from the time of concussion.*** The temporary exacerbation of mild symptoms with exercise is acceptable, as long as the symptoms quickly resolve at the completion of exercise, and as long as the exercise-related symptoms have completely resolved, before resumption of contact training.
- > The advanced care setting GRTSF is only available to adults aged 19 and above, who have consistent healthcare practitioner availability to monitor recovery throughout the return to sport/learn process.
- > If symptoms are severely exacerbated or remain for a prolonged period, review by a member of the clinical management team (CMT) or a health care practitioner (HCP) is recommended.
- > Several checkpoints are required to be passed to progress through the GRTSF and review by a healthcare practitioner is required prior to return to contact or high-risk activity.
- > Some athletes exposed to RHT develop CTE-NC. The factors that influence susceptibility to CTE-NC remain poorly understood.
- > ***If in doubt, sit them out.***

KEY POINTS FOR HEALTHCARE PRACTITIONERS

- > The diagnosis of concussion can be difficult. The signs and symptoms can be varied, non-specific and subtle. Athletes with suspected concussion should be removed from sport and assessed by a healthcare practitioner (HCP) of the clinical management team (CMT).
- > When assessing acute concussions, a standard initial assessment and management of the patient should be followed to ascertain whether there is any suggestion of neck or intracranial structural pathology.
- > Concussion is an evolving condition. Athletes suspected of, or diagnosed with, concussion require close monitoring and repeated assessment.
- > The diagnosis of concussion should be based on a clinical history and examination that includes a range of domains such as mechanism of injury, symptoms and signs, cognitive function, and neurology, including a balance assessment.
- > The **SCAT6** is currently the concussion assessment tool recommended internationally and covers the above-mentioned domains. It can be downloaded [here](#). **SCAT6** should not be used in isolation, but as part of the overall clinical assessment.
- > Computerised neurocognitive testing can be undertaken as part of the assessment but should not be used in isolation.
- > Blood tests are not indicated for uncomplicated concussion. Medical imaging is not indicated unless there is suspicion of more serious head, neck, or brain injury.
- > Standard head-injury advice should be given to all athletes suffering concussion and their carers.
- > Once the diagnosis of concussion has been made, immediate management is relative physical and cognitive rest. This includes time off school or work and deliberate rest from cognitive activity for 24–48 hours. After this period, the patient can return to light intensity physical activity that does not cause a significant and sustained deterioration in symptoms.
- > Children and adolescents, take longer to recover from concussion than adults.¹⁵⁴ A more conservative approach should be taken with those under the age of 19. ***The GRTSF requires those under 19 years of age and those without a dedicated HCP to guide recovery, to be symptom free for 14 days (at rest) before return to contact training, and not return to competitive contact sport until a minimum of 21 days from the time of concussion.*** To be clear, that is not 14 days from the time of concussion. It is 14 days from when the athlete becomes symptom-free. The day of the concussive incident is deemed day 0 of the GRTSF. The temporary exacerbation of mild symptoms with exercise is acceptable, as long as the symptoms quickly resolve at the completion of exercise, and as long as the exercise-related symptoms have completely resolved, before resumption of contact training.
- > An advanced care setting is only available for athletes with the fulfillment of certain eligibility criteria in the sporting organisation.
- > Some sports have their own guidelines or recommendations around the management of concussion in sport which should also be considered.
- > **If in doubt, sit them out.** *The athlete should not be returned to sport until cleared to do so by a health care practitioner with appropriate training and experience in concussion assessment and management.*
- > Where an athlete has had multiple concussions, the healthcare practitioner will counsel the athlete regarding the dangers of RHT, the potential long-term effects of RHT and the need for the athlete to think carefully about their continued involvement in high-risk sports, including contact and collision sports.
- > Some athletes that have been exposed to RHT may develop CTE-NC. The evidence regarding the association between RHT and CTE-NE consists of case reports, case series and retrospective analyses. The **degree of the association** between RHT and development of CTE-NC is unknown. The reliance on retired athletes nominating to posthumously undergo autopsy for this research generates significant bias in the samples examined. Confounding factors such as alcohol abuse, drug abuse, genetic predisposition, education exposure and psychiatric illness have not been controlled for adequately in studies conducted to date. Well-designed prospective studies are needed to better understand the possible relationship between RHT, concussion, and long-term brain health.

REFERENCES

1. Finch CF, Clapperton AJ, McCrory P. Increasing incidence of hospitalisation for sport-related concussion in Victoria, Australia. *Med J Aust.* 2013; 198(8):427–430.
2. Van Pelt KL, Puetz T, Swallow J, Lapointe AP, Broglio SP. Data-Driven Risk Classification of Concussion Rates: A Systematic Review and Meta-Analysis. *Sports Med.* 2021; 51(6):1227–1244.
3. Bretzin AC, Covassin T, Fox ME, et al. Sex differences in the clinical incidence of concussions, missed school days, and time loss in high school student-athletes: Part 1. *Am J Sports Med.* 2018; 46(9):2263–2269.
4. McCrory P, Meeuwisse W, Dvorak J, et al. Consensus statement on concussion in sport — the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med.* 2017; 51(11):838–847.
5. McCrory P, Feddermann-Demont N, Dvořák J, et al. What is the definition of sports-related concussion: a systematic review. *Br J Sports Med.* 2017; 51(11): 877–887.
6. Kamins J, Bigler E, Covassin T, et al. What is the physiological time to recovery after concussion? A systematic review. *Br J Sports Med.* 2017; 51(12): 935–940.
7. Weiler R, van Mechelen W, Fuller C, Ahmed OH, Verhagen E. Do neurocognitive SCAT3 baseline test scores differ between footballers (soccer) living with and without disability? A cross-sectional study. *Clin Sports Med.* 2018; 28(1):43–50.
8. Chandran A, Elmi A, Young H, DiPietro L. Determinants of concussion diagnosis, symptomology, and resolution time in U.S. high school soccer players. *Res Sports Med.* 2020; 28(1):42–54.
9. Black AM, Sergio LE, Macpherson AK. The epidemiology of concussions: number and nature of concussions and time to recovery among female and male canadian varsity athletes 2008 to 2011. *Clin J Sport Med.* 2017; 27(1):52–56.
10. Patricios JS, Schneider KJ, Dvorak J, et al. Consensus statement on concussion in sport: the 6th International Conference on Concussion in Sport—Amsterdam, October 2022. *Br J Sports Med.* 2023; 57(11):695–711.
11. Silverberg ND, Iverson GL, Cogan A, et al. The American Congress of Rehabilitation Medicine Diagnostic Criteria for Mild Traumatic Brain Injury. *Arch Phys Med Rehabil.* 2023; 104(8):1343–1355.
12. Feiss RS, Lutz M, Moody JR, Pangelinan MM. A systematic review of coach and parent knowledge of concussion. *J Concussion.* 2020; 4:2059700219900053.
13. Gourley MM, McLeod TCV, Bay RC. Awareness and recognition of concussion by youth athletes and their parents. *Athl. Train. Sports Health Care.* 2010; 2(5): 208–218.
14. Longworth T, McDonald A, Cunningham C, Khan H, Fitzpatrick J. Do rugby league players under-report concussion symptoms? A cross-sectional study of elite teams based in Australia. *BMJ Open Sport Exerc Med.* 2021; 7(1):e000860.
15. Meehan III W, Mannix RC, O'Brien MJ, Collins MW. The prevalence of undiagnosed concussions in athletes. *Clin J Sport Med.* 2013; 23(5):339–342.
16. Delaney JS, Caron JG, Correa JA, Bloom GA. Why professional football players chose not to reveal their concussion symptoms during a practice or game. *Clin Sports Med.* 2018; 28(1):1–12.
17. Koerte IK, Schultz V, Sydnor VJ, et al. Sex-related differences in the effects of sports-related concussion: a review. *J Neuroimaging.* 2020; 30(4):387–409.
18. Madsen TE, Bourjeily G, Hasnain M, et al. Sex-and gender-based medicine: the need for precise terminology. *Genet Genome.* 2017; 1(3):122–128.
19. National Rugby League. NRL injury surveillance report 2020 — executive summary 2020.
20. Gardner AJ, Iverson GL, Williams WH, Baker S, Stanwell P. A systematic review and meta-analysis of concussion in rugby union. *Sports Med.* 2014; 44(12):1717–1731.
21. Starling LT, Gabb N, Williams S, Kemp S, Stokes KA. Longitudinal study of six seasons of match injuries in elite female rugby union. *Br J Sports Med.* 2023; 57(4):212–217.
22. Australian Football League. 2021 AFL injury report. 2021.
23. Lian J, Sewani F, Dayan I, et al. Systematic Review of Injuries in the Men's and Women's National Basketball Association. *Am J Sports Med.* 2022; 50(5):1416–1429.
24. Fraas MR, Coughlan GF, Hart EC, McCarthy C. Concussion history and reporting rates in elite Irish rugby union players. *Phys Ther Sport.* 2014; 15(3): 136–142.

25. O'Connor S, Warrington G, Whelan G, McGoldrick A, Cullen S. Concussion history, reporting behaviors, attitudes, and knowledge in Jockeys. *Clin Sports Med*. 2020; 30(6).
26. Wallace J, Beidler E, Register-Mihalik JK, et al. Examining concussion nondisclosure in collegiate athletes using a health disparities framework and consideration of social determinants of health. *J Athl Train*. 2021; 57(1):16–24.
27. Australian Bureau of Statistics (ABS). Estimates of Aboriginal and Torres Strait Islander Australians [Internet]. Canberra: ABS; 2021 June [cited 2023 January 19]. Available from: <https://www.abs.gov.au/statistics/people/aboriginal-and-torres-strait-islander-peoples/estimates-aboriginal-and-torres-strait-islander-australians/latest-release>.
28. Australian Rugby League Commission. Submission to the House of Representatives Standing Committee on Aboriginal and Torres Strait Islander Affairs inquiry into the contribution of sport to Indigenous wellbeing and mentoring. Canberra 2012.
29. Lakhani A, Townsend C, Bishara J. Traumatic brain injury amongst indigenous people: a systematic review. *Brain Injury*. 2017; 31(13–14):1718–1730.
30. Wright M, Brown A, Dudgeon P, et al. Our journey, our story: a study protocol for the evaluation of a co-design framework to improve services for Aboriginal youth mental health and well-being. *BMJ Open*. 2021; 11(5):e042981.
31. Abrahams S, Fie SM, Patricios J, Posthumus M, September AV. Risk factors for sports concussion: an evidence-based systematic review. *Br J Sports Med*. 2014; 48(2):91–97.
32. McGroarty NK, Brown SM, Mulcahey MK. Sport-related concussion in female athletes: a systematic review. *Orthop J Sports Med*. 2020; 8(7):2325967120932306–2325967120932306.
33. Covassin T, Savage JL, Bretzin AC, Fox ME. Sex differences in sport-related concussion long-term outcomes. *Int J Psychophysiol*. 2018; 132:9–13.
34. Lincoln AE, Caswell SV, Almquist JL, Dunn RE, Norris JB, Hinton RY. Trends in concussion incidence in high school sports: a prospective 11-year study. *Am J Sports Med*. 2011; 39(5):958–963.
35. Marar M, McIlvain NM, Fields SK, Comstock RD. Epidemiology of concussions among United States high school athletes in 20 sports. *Am J Sports Med*. 2012; 40(4):747–755.
36. Zuckerman SL, Kerr ZY, Yengo-Kahn A, Wasserman E, Covassin T, Solomon GS. Epidemiology of sports-related concussion in NCAA athletes from 2009–2010 to 2013–2014: incidence, recurrence, and mechanisms. *Am J Sports Med*. 2015; 43(11):2654–2662.
37. Gardner A, Kay-Lambkin F, Stanwell P, et al. A systematic review of diffusion tensor imaging findings in sports-related concussion. *J Neurotrauma*. 2012; 29(16):2521–2538.
38. Tierney RT, Sitler MR, Swanik CB, Swanik KA, Higgins M, Torg J. Gender differences in head-neck segment dynamic stabilization during head acceleration. *Med Sci Sports Exerc*. 2005; 37(2):272–279.
39. Hootman JM, Dick R, Agel J. Epidemiology of collegiate injuries for 15 sports: summary and recommendations for injury prevention initiatives. *J Athl Train*. 2007; 42(2):311–319.
40. Chandran A, Boltz AJ, Morris SN, et al. Epidemiology of concussions in National Collegiate Athletic Association (NCAA) sports: 2014/15–2018/19. *Am J Sports Med*. 2022; 50(2):526–536.
41. Williams EMP, Petrie FJ, Pennington TN, et al. Sex differences in neck strength and head impact kinematics in university rugby union players. *Eur J Sport Sci*. 2021:1–10.
42. Wilcox BJ, Beckwith JG, Greenwald RM, et al. Biomechanics of head impacts associated with diagnosed concussion in female collegiate ice hockey players. *J Biomech*. 2015; 48(10):2201–2204.
43. Guskiewicz KM, Mihalik JP, Shankar V, et al. Measurement of head impacts in collegiate football players: relationship between head impact biomechanics and acute clinical outcome after concussion. *Neurosurgery*. 2007; 61(6):1244–1253.
44. Lynall RC, Clark MD, Grand EE, et al. Head impact biomechanics in women's college soccer. *Med Sci Sports Exerc*. 2016; 48(9):1772–1778.
45. Bretzin AC, Covassin T, Wiebe DJ, Stewart W. Association of sex with adolescent soccer concussion incidence and characteristics. *JAMA Netw Open*. 2021; 4(4):e218191.
46. Di Battista AP, Rhind SG, Churchill N, Richards D, Lawrence DW, Hutchison MG. Peripheral blood neuroendocrine hormones are associated with clinical indices of sport-related concussion. *Scientific reports*. 2019; 9(1):18605.
47. La Fountaine MF, Hill-Lombardi V, Hohn AN, Leahy CL, Testa AJ. Preliminary evidence for a window of increased vulnerability to sustain a concussion in females: a brief report. *Front Neurol*. 2019; 10.

48. Wunderle K, Hoeger KM, Wasserman E, Bazarian JJ. Menstrual phase as predictor of outcome after mild traumatic brain injury in women. *J Head Trauma Rehabil.* 2014; 29(5):E1–8.
49. Snook ML, Henry LC, Sanfilippo JS, Zeleznik AJ, Kontos AP. Association of concussion with abnormal menstrual patterns in adolescent and young women. *JAMA Pediatr.* 2017; 171(9):879–886.
50. Wagner J, Dusick JR, McArthur DL, et al. Acute gonadotroph and somatotroph hormonal suppression after traumatic brain injury. *J Neurotrauma.* 2010; 27(6):1007–1019.
51. Pieroth EM, Wicklund A. Concussion in Female Athletes. *Operative Techniques in Sports Medicine.* 2023:151026.
52. Miyashita TL, Diakogeorgiou E, VanderVegt C. Gender Differences in Concussion Reporting Among High School Athletes. *Sports Health.* 2016; 8(4):359–363.
53. Tweedy SM, Vanlandewijck YC. International Paralympic Committee position stand-background and scientific principles of classification in Paralympic sport. *Br J Sports Med.* 2011; 45(4):259–269.
54. Kissick J, Webborn N. Concussion in para sport. *Phys Med Rehabil Clin N Am.* 2018; 29(2):299–311.
55. Webborn N, Cushman D, Blauwet CA, et al. The Epidemiology of Injuries in Football at the London 2012 Paralympic Games. *PM R.* 2016; 8(6):545–552.
56. Derman W, Runciman P, Schwellnus M, et al. High precompetition injury rate dominates the injury profile at the Rio 2016 Summer Paralympic Games: a prospective cohort study of 51198 athlete days. *Br J Sports Med.* 2018; 52(1):24–31.
57. Derman W, Runciman P, Jordaan E, et al. High incidence of injuries at the Pyeongchang 2018 Paralympic Winter Games: a prospective cohort study of 6804 athlete days. *Br J Sports Med.* 2020; 54(1):38–43.
58. Centers for Disease Control and Prevention. CDC announces updated information to help physicians recognize and manage concussions early 2007.
59. Arbogast KB, Curry AE, Pfeiffer MR, et al. Point of health care entry for youth with concussion within a large pediatric care network. *JAMA pediatrics.* 2016; 170(7):e160294.
60. Karlin AM. Concussion in the pediatric and adolescent population: "different population, different concerns". *PM R.* 2011; 3(10 Suppl 2):S369–379.
61. Browne G. DS. Concussive head injury in children and adolescents. *Aust Fam Physician.* 2016; 45(7):470–476.
62. Browne GJ, Lam LT. Concussive head injury in children and adolescents related to sports and other leisure physical activities. *Br J Sports Med.* 2006; 40(2):163–168.
63. Chun BJ, Furutani T, Oshiro R, Young C, Prentiss G, Murata N. Concussion Epidemiology in Youth Sports: Sports Study of a Statewide High School Sports Program. *Sports Health.* 2021; 13(1):18–24.
64. Pfister T, Pfister K, Hagel B, Ghali WA, Ronksley PE. The incidence of concussion in youth sports: a systematic review and meta-analysis. *Br J Sports Med.* 2016; 50(5):292.
65. Warren D. Childrens use of health care services. In Growing up in Australia- The Longitudinal Study of Australian Children, Annual Statistical Report 2017. *Growing Up in Australia.* 2018.
66. Arain M, Haque M, Johal L, et al. Maturation of the adolescent brain. *Neuropsychiatr Dis Treat.* 2013; 9:449–461.
67. Giedd JN. Structural magnetic resonance imaging of the adolescent brain. *Ann N Y Acad Sci.* 2004; 1021:77–85.
68. Giedd JN, Blumenthal J, Jeffries NO, et al. Brain development during childhood and adolescence: a longitudinal MRI study. *Nat Neurosci.* 1999; 2(10):861–863.
69. Australian Insititute of Health and Welfare (AIHW). Hospital separations due to traumatic brain injury, Australia 2004–05. Canberra: *AIHW*; 2008.
70. Thomas E, Fitzgerald M, Cowen G. Does Australia have a concussion 'epidemic'? *Concussion.* 2020; 5(1):Cnc70.
71. Steenerson K, Starling AJ. Pathophysiology of sports-related concussion. *Neurologic clinics.* 2017; 35(3):403–408.
72. Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery.* 2014; 75 Suppl 4:S24–33.
73. Martin G. Traumatic brain injury: The first 15 milliseconds. *Brain Inj.* 2016; 30(13–14):1517–1524.
74. Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery.* 2014; 75 Suppl 4(0 4):S24–33.
75. Nopoulos P, Flaum M, O'Leary D, Andreasen NC. Sexual dimorphism in the human brain: evaluation of tissue volume, tissue composition and surface anatomy using magnetic resonance imaging. *Psychiatry Res.* 2000; 98(1):1–13.

76. Dollé JP, Jaye A, Anderson SA, Ahmadzadeh H, Shenoy VB, Smith DH. Newfound sex differences in axonal structure underlie differential outcomes from in vitro traumatic axonal injury. *Exp Neurol*. 2018; 300:121–134.
77. Chen JK, Johnston KM, Petrides M, Ptito A. Recovery from mild head injury in sports: evidence from serial functional magnetic resonance imaging studies in male athletes. *Clin J Sport Med*. 2008; 18(3):241–247.
78. Palacios EM, Yuh EL, Mac Donald CL, et al. Diffusion tensor imaging reveals elevated diffusivity of white matter microstructure that is independently associated with long-term outcome after mild traumatic brain injury: a TRACK-TBI study. *J Neurotrauma*. 2022; 39(19–20):1318–1328.
79. Barlow KM, Marcil LD, Dewey D, et al. Cerebral perfusion changes in post-concussion syndrome: a prospective controlled cohort study. *J Neurotrauma*. 2016; 34(5):996–1004.
80. Prichep LS, McCrear M, Barr W, Powell M, Chabot RJ. Time course of clinical and electrophysiological recovery after sport-related concussion. *J Head Trauma Rehabil*. 2013; 28(4):266–273.
81. Peskind ER, Petrie EC, Cross DJ, et al. Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms. *Neuroimage*. 2011; 54 Suppl 1:S76–82.
82. Eliason PH, Galarneau J-M, Kolstad AT, et al. Prevention strategies and modifiable risk factors for sport-related concussions and head impacts: a systematic review and meta-analysis. *Br J Sports Med*. 2023; 57(12):749–761.
83. Pankow MP, Syrydiuk RA, Kolstad AT, et al. Head Games: A Systematic Review and Meta-analysis Examining Concussion and Head Impact Incidence Rates, Modifiable Risk Factors, and Prevention Strategies in Youth Tackle Football. *Sports Med*. 2022; 52(6):1259–1272.
84. Guillaume S, Lincoln AE, Hepburn L, Caswell SV, Kerr ZY. Rule Modifications to Reduce Checking-Related Injuries in High School Boys' Lacrosse. *J Athl Train*. 2021; 56(4):437–445.
85. MEDICINE COS, FITNESS, Brenner JS, et al. Tackling in Youth Football. *Pediatrics*. 2015; 136(5):e1419–e1430.
86. Shanley E, Thigpen C, Kissenberth M, et al. Heads Up Football Training Decreases Concussion Rates in High School Football Players. *Clin J Sport Med*. 2021; 31(2):120–126.
87. Kerr ZY, Yeargin S, Valovich McLeod TC, et al. Comprehensive Coach Education and Practice Contact Restriction Guidelines Result in Lower Injury Rates in Youth American Football. *Orthop J Sports Med*. 2015; 3(7):2325967115594578.
88. Rugby.com. Tackle height trial: What it really means for Rugby. Accessed 2023 13 December. From [Tackle height trial: What it really means for Rugby](#)
89. Rugby-League.com. Making Rugby League safer and more accessible — significant changes to be introduced at all levels from 2024. Accessed 2023 13 December. From [Making Rugby League safer and more accessible — significant changes to be introduced at all levels from 2024 \(rugby-league.com\)](#)
90. Daneshvar DH, Nowinski CJ, McKee AC, Cantu RC. The epidemiology of sport-related concussion. *Clin Sports Med*. 2011; 30(1):1–17, vii.
91. Hoshizaki TB, Post A, Oeur RA, Brien SE. Current and future concepts in helmet and sports injury prevention. *Neurosurgery*. 2014; 75 Suppl 4:S136–148.
92. Makovec Knight J, Nguyen JVK, Mitra B, Willmott C. Soft-shell headgear, concussion and injury prevention in youth team collision sports: a systematic review. *BMJ Open*. 2021; 11(6):e044320.
93. van Ierssel J, Ledoux AA, Tang K, Zemek R. Sex-based differences in symptoms with mouthguard use after pediatric sport-related concussion. *J Athl Train*. 2021; 56(11):1188–1196.
94. Greenhill DA, Navo P, Zhao H, Torg J, Comstock RD, Boden BP. Inadequate Helmet Fit Increases Concussion Severity in American High School Football Players. *Sports Health*. 2016; 8(3):238–243.
95. Cecchi NJ, Domel AG, Liu Y, et al. Identifying Factors Associated with Head Impact Kinematics and Brain Strain in High School American Football via Instrumented Mouthguards. *Ann Biomed Eng*. 2021; 49(10):2814–2826.
96. McGuine TA, Hetzel S, McCrear M, Brooks MA. Protective Equipment and Player Characteristics Associated With the Incidence of Sport-Related Concussion in High School Football Players: A Multifactorial Prospective Study. *Am J Sports Med*. 2014; 42(10):2470–2478.
97. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J. Examining Concussion Rates and Return to Play in High School Football Players Wearing Newer Helmet Technology: A Three-Year Prospective Cohort Study. *Neurosurgery*. 2006; 58(2):275–286.

98. Delaney JS, Al-Kashmiri A, Drummond R, Correa JA. The effect of protective headgear on head injuries and concussions in adolescent football (soccer) players. *Br J Sports Med.* 2008; 42(2):110–115.
99. McGuine T, Post E, Pfaller AY, et al. Does soccer headgear reduce the incidence of sport-related concussion? A cluster, randomised controlled trial of adolescent athletes. *Br J Sports Med.* 2020; 54(7): 408–413.
100. Collins CL, Fletcher EN, Fields SK, et al. Neck strength: a protective factor reducing risk for concussion in high school sports. *J Prim Prev.* 2014; 35(5):309–319.
101. Waring KM, Smith ER, Austin GP, Bowman TG. Exploring the effects of a neck strengthening program on purposeful soccer heading biomechanics and neurocognition. *Int J Sports Phys Ther.* 2022; 17(6):1043–1052.
102. Cooney NJ, Sowman P, Schilaty N, Bates N, Hewett TE, Doyle TLA. Head and Neck Characteristics as Risk Factors For and Protective Factors Against Mild Traumatic Brain Injury in Military and Sporting Populations: A Systematic Review. *Sports Med.* 2022; 52(9):2221–2245.
103. Garcia GP, Czerniak LL, Lavieri MS, et al. Estimating the Relationship Between the Symptom-Free Waiting Period and Injury Rates After Return-to-Play from Concussion: A Simulation Analysis Using CARE Consortium Data. *Sports Med.* 2023.
104. Eliason PH, Galarneau J-M, Kolstad AT, et al. Prevention strategies and modifiable risk factors for sport-related concussions and head impacts: a systematic review and meta-analysis. *B J Sports Med.* 2023; 57(12):749–761.
105. Patricios JS, Arden CL, Hislop MD, et al. Implementation of the 2017 Berlin Concussion in Sport Group Consensus Statement in contact and collision sports: a joint position statement from 11 national and international sports organisations. *Br J Sports Med.* 2018; 52(10):635–641.
106. Echemendia RJ, Burma JS, Bruce JM, et al. Acute evaluation of sport-related concussion and implications for the Sport Concussion Assessment Tool (SCAT6) for adults, adolescents and children: a systematic review. *Br J Sports Med.* 2023; 57(11):722–735.
107. Brown L, Camarinos J. The Role of Physical Therapy in Concussion Rehabilitation. *Semin Pediatr Neurol.* 2019; 30:68–78.
108. Makdissi M, Cantu RC, Johnston KM, McCrory P, Meeuwisse WH. The difficult concussion patient: what is the best approach to investigation and management of persistent (>10 days) postconcussive symptoms? *Br J Sports Med.* 2013; 47(5):308–313.
109. Collins MW, Kontos AP, Reynolds E, Murawski CD, Fu FH. A comprehensive, targeted approach to the clinical care of athletes following sport-related concussion. *Knee Surg Sports Traumatol Arthrosc.* 2014; 22(2):235–246.
110. Lyons TW, Mannix R, Tang K, et al. Paediatric post-concussive symptoms: symptom clusters and clinical phenotypes. *Br J Sports Med.* 2022; 56(14):785–791.
111. Davis G, Rausa V, Babl F, et al. Improving subacute management of post concussion symptoms: a pilot study of the Melbourne Paediatric Concussion Scale parent report. *Concussion.* 2022; 7.
112. Quatman-Yates CC, Hunter-Giordano A, Shimamura KK, et al. Physical therapy evaluation and treatment after concussion/mild traumatic brain injury. *J Orthop Sports Phys Ther.* 2020; 50(4):CPG1–CPG73.
113. Kontos AP, Sufrinko A, Sandel N, Emami K, Collins MW. Sport-related Concussion Clinical Profiles: Clinical Characteristics, Targeted Treatments, and Preliminary Evidence. *Curr Sports Med Rep.* 2019; 18(3):82–92.
114. Anzalone AJ, Blueitt D, Case T, et al. A positive vestibular/ocular motor screening (VOMS) is associated with increased recovery time after sports-related concussion in youth and adolescent athletes. *Am J Sports Med.* 2017; 45(2):474–479.
115. Ellis MJ, Cordingley DM, Vis S, Reimer KM, Leiter J, Russell K. Clinical predictors of vestibulo-ocular dysfunction in pediatric sports-related concussion. *J Neurosurg Pediatr.* 2017; 19(1):38–45.
116. Gray M, Wilson JC, Potter M, Provance AJ, Howell DR. Female adolescents demonstrate greater oculomotor and vestibular dysfunction than male adolescents following concussion. *Phys Ther Sport.* 2020; 42:68–74.
117. Leddy JJ, Baker JG, Willer B. Active rehabilitation of concussion and post-concussion syndrome. *Phys Med Rehabil Clin N Am.* 2016; 27(2):437–454.
118. Iverson GL, Gardner AJ, Terry DP, et al. Predictors of clinical recovery from concussion: a systematic review. *Br J Sports Med.* 2017; 51(12):941–948.
119. Moran RN, Wallace J, Murray NG, Covassin T. Effects of attention deficit hyperactivity disorder and learning disability on vestibular and ocular baseline concussion assessment in pediatric athletes. *Appl Neuropsychol Child.* 2021; 10(3):276–282.

120. King DA, Hume PA, Hind K, Clark TN, Hardaker N. The Incidence, Cost, and Burden of Concussion in Women's Rugby League and Rugby Union: A Systematic Review and Pooled Analysis. *Sports Med.* 2022; 52(8):1751–1764.
121. Frommer LJ, Gurka KK, Cross KM, Ingersoll CD, Comstock RD, Saliba SA. Sex differences in concussion symptoms of high school athletes. *J Athl Train.* 2011; 46(1):76–84.
122. Merritt VC, Arnett PA. Premorbid predictors of postconcussion symptoms in collegiate athletes. *J Clin Exp Neuropsychol.* 2014; 36(10):1098–1111.
123. Ono KE, Burns TG, Bearden DJ, McManus SM, King H, Reisner A. Sex-based differences as a predictor of recovery trajectories in young athletes after a sports-related concussion. *Am J Sports Med.* 2016; 44(3):748–752.
124. Preiss-Farzanegan SJ, Chapman B, Wong TM, Wu J, Bazarian JJ. The relationship between gender and postconcussion symptoms after sport-related mild traumatic brain injury. *PM R.* 2009; 1(3):245–253.
125. Covassin T, Elbin RJ, Bleecker A, Lipchik A, Kontos AP. Are there differences in neurocognitive function and symptoms between male and female soccer players after concussions? *Am J Sports Med.* 2013; 41(12):2890–2895.
126. Covassin T, Elbin RJ, Harris W, Parker T, Kontos A. The role of age and sex in symptoms, neurocognitive performance, and postural stability in athletes after concussion. *Am J Sports Med.* 2012; 40(6):1303–1312.
127. Covassin T, Schatz P, Swanik CB. Sex differences in neuropsychological function and post-concussion symptoms of concussed collegiate athletes. *Neurosurgery.* 2007; 61(2):345–350; discussion 350–341.
128. Brown DA, Elsass JA, Miller AJ, Reed LE, Reneker JC. Differences in Symptom Reporting Between Males and Females at Baseline and After a Sports-Related Concussion: A Systematic Review and Meta-Analysis. *Sports Med.* 2015; 45(7):1027–1040.
129. Baker JG, Leddy JJ, Darling SR, Shucard J, Makdissi M, Willer BS. Gender differences in recovery from sports-related concussion in adolescents. *Clin Pediatr.* 2016; 55(8):771–775.
130. Al-Hassany L, Haas J, Piccininni M, Kurth T, Maassen Van Den Brink A, Rohmann JL. Giving researchers a headache — sex and gender differences in migraine. *Front Neurol.* 2020; 11.
131. O'Connor KL, Dain Allred C, Cameron KL, et al. Descriptive analysis of a baseline concussion battery among U.S. Service Academy members: results from the Concussion Assessment, Research, and Education (CARE) consortium. *Mil Med.* 2018; 183(11–12):e580–e590.
132. McCrea M, Pritchep L, Powell MR, Chabot R, Barr WB. Acute effects and recovery after sport-related concussion: a neurocognitive and quantitative brain electrical activity study. *J Head Trauma Rehabil.* 2010; 25(4):283–292.
133. Maddocks DL, Dicker GD, Saling MM. The assessment of orientation following concussion in athletes. *Clin J Sport Med.* 1995; 5(1):32–35.
134. Patricios J, Fuller GW, Ellenbogen R, et al. What are the critical elements of sideline screening that can be used to establish the diagnosis of concussion? A systematic review. *Br J Sports Med.* 2017; 51(11):888–894.
135. Sport Concussion Assessment Tool 6 [SCAT6]. *Br J Sports Med.* 2023; 57(11):622–631.
136. Echemendia RJ, Meeuwisse W, McCrory P, et al. The sport concussion assessment tool 5th edition [SCAT5]: background and rationale. *Br J Sports Med.* 2017; 51(11):848–850.
137. Bruce JM, Echemendia RJ, Meeuwisse W, Hutchison MG, Aubry M, Comper P. Development of a risk prediction model among professional hockey players with visible signs of concussion. *Br J Sports Med.* 2018; 52(17):1143–1148.
138. Downey RI, Hutchison MG, Comper P. Determining sensitivity and specificity of the Sport Concussion Assessment Tool 3 [SCAT3] components in university athletes. *Brain Inj.* 2018; 32(11):1345–1352.
139. Echemendia RJ, Thelen J, Meeuwisse W, Comper P, Hutchison MG, Bruce JM. Testing the hybrid battery approach to evaluating sports-related concussion in the National Hockey League: A factor analytic study. *Clin Neuropsychol.* 2020; 34(5):899–918.
140. Echemendia RJ, Broglio SP, Davis GA, et al. What tests and measures should be added to the SCAT3 and related tests to improve their reliability, sensitivity and/or specificity in sideline concussion diagnosis? A systematic review. *Br J Sports Med.* 2017; 51(11):895–901.
141. Bullen J, Hill-Wall T, Norman R, Cowen G. Concussion in Aboriginal and Torres Strait Islander peoples: what is the true epidemiology? *J Neurotrauma.* 2021; 38:1411–1440.

142. Holtzhausen LJ, Souissi S, Sayrafi OA, et al. Arabic translation and cross-cultural adaptation of the Sport Concussion Assessment Tool 5 (SCAT5). *Biol Sport*. 2021; 38(1):129–144.
143. The Concussion Recognition Tool 6 (CRT6). *Br J Sports Med*. 2023; 57(11):692–694.
144. Weiler R, Blauwet C, Clarke D, et al. Concussion in para sport: the first position statement of the Concussion in Para Sport (CIPS) Group. *Br J Sports Med*. 2021; 55(21):1187–1195.
145. Webborn N, Blauwet CA, Derman W, et al. Heads up on concussion in para sport. *Br J Sports Med*. 2018; 52(18):1157–1158.
146. West LR, Griffin S, Weiler R, Ahmed OH. Management of concussion in disability sport: a different ball game? *Br J Sports Med*. 2017; 51(14):1050–1051.
147. Griffin S, West LR, Ahmed OH, Weiler R. Concussion knowledge, attitudes, and beliefs amongst sports medicine personnel at the 2015 cerebral palsy football world championships. *Br J Sports Med*. 2017; 51(4):325–325.
148. Wright MJ, Woo E, Birath JB, et al. An index predictive of cognitive outcome in retired professional American Football players with a history of sports concussion. *J Clin Exp Neuropsychol*. 2016; 38(5):561–571.
149. Echemendia RJ, Meeuwisse W, McCrory P, et al. The concussion recognition tool 5th edition (CRT5): background and rationale. *Br J Sports Med*. 2017; 51(11):870–871.
150. Concussion recognition tool 5(c). *Br J Sports Med*. 2017; 51(11):872.
151. Sport Concussion Office Assessment Tool 6 (SCOAT6). *Br J Sports Med*. 2023; 57(11):651–667.
152. Patricios JS, Davis GA, Ahmed OH, et al. Introducing the Sport Concussion Office Assessment Tool 6 (SCOAT6). *Br J Sports Med*. 2023; 57(11):648–650.
153. Sport concussion assessment tool for childrens ages 5 to 12 years. *Br J Sports Med*. 2017; 51(11):862–869.
154. Davis GA, Echemendia RJ, Ahmed OH, et al. Introducing the Child Sport Concussion Assessment Tool 6 (Child SCAT6). *Br J Sports Med*. 2023; 57(11):632–635.
155. Echemendia R, Burma J, Bruce J. Acute detection of sport-related concussion and implications for the SCAT tools: A systematic review. *Br J Sports Med*. 2023; 57:722–735.
156. Davis GA, Patricios JS, Purcell LK, et al. Introducing the Child Sport Concussion Office Assessment Tool 6 (Child SCOAT6). *Br J Sports Med*. 2023; 57(11):668–671.
157. Child SCOAT6. *Br J Sports Med*. 2023; 57(11):672–688.
158. Davis GA, Rausa VC, Babl FE, et al. Improving subacute management of post concussion symptoms: a pilot study of the Melbourne Paediatric Concussion Scale parent report. *Concussion*. 2022; 7(1):CNC97.
159. Ayr LK, Yeates KO, Taylor HG, Browne M. Dimensions of postconcussive symptoms in children with mild traumatic brain injuries. *J Int Neuropsychol Soc*. 2009; 15(1):19–30.
160. Pearson R, Sheridan CA, Kang K, et al. Post-concussive orthostatic tachycardia is distinct from postural orthostatic tachycardia syndrome (POTS) in children and adolescents. *Child Neurology Open*. 2022; 9:2329048X221082753.
161. Corwin DJ, Arbogast KB, Swann C, Haber R, Grady MF, Master CL. Reliability of the visio-vestibular examination for concussion among providers in a pediatric emergency department. *Am J Emerg Med*. 2020; 38(9):1847–1853.
162. Ramsey KA, Vaughan C, Wagner BM, McGuire JF, Gioia GA. Impact of Self-Efficacy and Affective Functioning on Pediatric Concussion Symptom Severity. *J Int Neuropsychol Soc*. 2021; 27(9):875–882.
163. Uchiyama CL, D'Elia LF, Dellinger AM, et al. Longitudinal comparison of alternate versions of the symbol digit modalities test: Issues of form comparability and moderating demographic variables. *Clin Neuropsychol*. 1994; 8(2):209–218.
164. Pilkonis PA, Choi SW, Reise SP, Stover AM, Riley WT, Cella D. Item banks for measuring emotional distress from the Patient-Reported Outcomes Measurement Information System (PROMIS®): depression, anxiety, and anger. *Assessment*. 2011; 18(3):263–283.
165. Irwin DE, Stucky B, Langer MM, et al. An item response analysis of the pediatric PROMIS anxiety and depressive symptoms scales. *Qual Life Res*. 2010; 19(4):595–607.
166. Cairncross M, Brooks BL, Virani S, Silverberg ND. Fear avoidance behavior in youth with poor recovery from concussion: measurement properties and correlates of a new scale. *Child Neuropsychol*. 2021; 27(7):911–921.
167. Broglio SP, Collins MW, Williams RM, Mucha A, Kontos AP. Current and emerging rehabilitation for concussion: a review of the evidence. *Clin Sports Med*. 2015; 34(2):213–231.
168. Kontos AP, Deitrick JM, Collins MW, Mucha A. Review of vestibular and oculomotor screening and concussion rehabilitation. *J Athl Train*. 2017; 52(3):256–261.

169. Lumba-Brown A, Teramoto M, Bloom OJ, et al. Concussion guidelines step 2: evidence for subtype classification. *Neurosurgery*. 2020; 86(1):2–13.
170. McLeod TC, Lewis JH, Whelihan K, Bacon CE. Rest and return to activity after sport-related concussion: a systematic review of the literature. *J Athl Train*. 2017; 52(3):262–287.
171. Thomas DG, Apps JN, Hoffmann RG, McCrea M, Hammeke T. Benefits of strict rest after acute concussion: a randomized controlled trial. *Pediatrics*. 2015; 135(2):213–223.
172. Macnow T, Curran T, Tolliday C, et al. Effect of screen time on recovery from concussion: a randomized clinical trial. *JAMA Pediatr*. 2021; 175(11):1124–1131.
173. Wiebe DJ, Bretzin AC, D'Alonzo BA. Progression through return-to-sport and return-to-academics guidelines for concussion management and recovery in collegiate student athletes: findings from the Ivy League-Big Ten Epidemiology of Concussion Study. *Br J Sports Med*. 2022; 56(14):801–811.
174. Worts PR, Mason JR, Burkhart SO, Sanchez-Gonzalez MA, Kim JS. The acute, systemic effects of aerobic exercise in recently concussed adolescent student-athletes: preliminary findings. *Eur J Appl Physiol*. 2022; 122(6):1441–1457.
175. Schneider KJ, Leddy JJ, Guskiewicz KM, et al. Rest and treatment/rehabilitation following sport-related concussion: a systematic review. *Br J Sports Med*. 2017; 51(12):930–934.
176. Reid SA, Farbenblum J, McLeod S. Do physical interventions improve outcomes following concussion: a systematic review and meta-analysis? *Br J Sports Med*. 2022; 56(5):292.
177. McDevitt J, Appiah-Kubi KO, Tierney R, Wright WG. Vestibular and oculomotor assessments may increase accuracy of subacute concussion assessment. *Int J Sports Med*. 2016; 37(9):738–747.
178. Pryhoda MK, Shelburne KB, Gorgens K, Ledreux A, Granholm AC, Davidson BS. Centre of pressure velocity shows impairments in NCAA Division I athletes six months post-concussion during standing balance. *J Sports Sci*. 2020; 38(23):2677–2687.
179. Sherry NS, Fazio-Sumrok V, Sufrinko A, Collins MW, Kontos AP. Multimodal assessment of sport-related concussion. *Clin Sports Med*. 2021; 31(3):244–249.
180. Cheever KM, McDevitt J, Tierney R, Wright WG. Concussion Recovery phase affects vestibular and oculomotor symptom provocation. *Int J Sports Med*. 2018; 39(2):141–147.
181. Cohen PE, Sufrinko A, Elbin RJ, Collins MW, Sinnott AM, Kontos AP. Do initial symptom factor scores predict subsequent impairment following concussion? *Clin J Sport Med*. 2020; 30 Suppl 1(Suppl 1):S61–s68.
182. Eagle SR, Kontos AP, Sinnott A, et al. Utility of a novel perceptual-motor control test for identification of sport-related concussion beyond current clinical assessments. *J Sports Sci*. 2020; 38(15):1799–1805.
183. Elbin RJ, Sufrinko A, Anderson MN, et al. Prospective changes in vestibular and ocular motor impairment after concussion. *J Neurol Phys Ther*. 2018; 42(3):142–148.
184. Kontos AP, Elbin RJ, Trbovich A, et al. Concussion clinical profiles screening (CP Screen) tool: preliminary evidence to inform a multidisciplinary approach. *Neurosurgery*. 2020; 87(2):348–356.
185. Leddy J, Lesh K, Haider MN, et al. Derivation of a focused, brief concussion physical examination for adolescents with sport-related concussion. *Clin J Sport Med*. 2021; 31(1):7–14.
186. Mucha A, Collins MW, Elbin RJ, et al. A brief vestibular/ocular motor screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med*. 2014; 42(10):2479–2486.
187. Sufrinko A, McAllister-Deitrick J, Elbin RJ, Collins MW, Kontos AP. Family history of migraine associated with posttraumatic migraine symptoms following sport-related concussion. *J Head Trauma Rehabil*. 2018; 33(1):7–14.
188. Sufrinko AM, Howie EK, Charek DB, Elbin RJ, Collins MW, Kontos AP. Mobile ecological momentary assessment of postconcussion symptoms and recovery outcomes. *J Head Trauma Rehabil*. 2019; 34(6):E40–e48.
189. Vernau BT, Grady MF, Goodman A, et al. Oculomotor and neurocognitive assessment of youth ice hockey players: baseline associations and observations after concussion. *Dev Neuropsychol*. 2015; 40(1):7–11.
190. Sufrinko AM, Cohen PE, Kontos AP, Marchetti GF, Elbin RJ, Re V. Using acute performance on a comprehensive neurocognitive, vestibular, and ocular motor assessment battery to predict recovery duration after sport-related concussions. *Am J Sports Med*. 2017; 45(5):1187–1194.
191. Price AM, Knell G, Caze TJ, 2nd, Abt JP, Loveland D, Burkhart SO. Exploring Vestibular/Ocular and Cognitive Dysfunction as Prognostic Factors for Prolonged Recovery in Sports-Related Concussion Patients Aged 8 to 12 Years. *Clin J Sport Med*. 2022; 32(4):408–414.

192. Van Dyke SA, Axelrod BN, Schutte C. The utility of the post-concussive symptom questionnaire. *Arch Clin Neuropsychol*. 2010; 25(7):634–639.
193. Marshall S, Bayley M, McCullagh S, et al. Updated clinical practice guidelines for concussion/mild traumatic brain injury and persistent symptoms. *Brain Inj*. 2015; 29(6):688–700.
194. Grool AM, Aglipay M, Momoli F, et al. Association between early participation in physical activity following acute concussion and persistent postconcussive symptoms in children and adolescents. *JAMA*. 2016; 316(23):2504–2514.
195. Wang EX, Hwang CE, Nguyen JN, Segovia NA, Abrams GD, Kussman A. Factors Associated With a Prolonged Time to Return to Play After a Concussion. *Am J Sports Med*. 2022; 50(6):1695–1701.
196. Makdissi M, Critchley ML, Cantu RC, et al. When should an athlete retire or discontinue participating in contact or collision sports following sport-related concussion? A systematic review. *Br J Sports Med*. 2023; 57(12):822–830.
197. Di Pietro V, O'Halloran P, Watson CN, et al. Unique diagnostic signatures of concussion in the saliva of male athletes: the Study of Concussion in Rugby Union through MicroRNAs (SCRUM). *Br J Sports Med*. 2021; 55(24):1395–1404.
198. Hicks SD, Onks C, Kim RY, et al. Refinement of saliva microRNA biomarkers for sports-related concussion. *J Sport Health Sci*. 2021.
199. McCrea M, Broglio SP, McAllister TW, et al. Association of blood biomarkers with acute sport-related concussion in collegiate athletes: findings from the NCAA and Department of Defense CARE consortium. *JAMA Network Open*. 2020; 3(1):e1919771–e1919771.
200. Mitra B, Rau TF, Surendran N, et al. Plasma micro-RNA biomarkers for diagnosis and prognosis after traumatic brain injury: A pilot study. *J Clin Neurosci*. 2017; 38:37–42.
201. Shultz SR, Taylor CJ, Aggio-Bruce R, et al. Decrease in plasma miR-27a and miR-221 after concussion in Australian Football players. *Biomark Insights*. 2022; 17:11772719221081318.
202. Mitra B, Reyes J, O'Brien WT, et al. Micro-RNA levels and symptom profile after mild traumatic brain injury: a longitudinal cohort study. *J Clin Neurosci*. 2022; 95:81–87.
203. Manley G, Gardner AJ, Schneider KJ, et al. A systematic review of potential long-term effects of sport-related concussion. *Br J Sports Med*. 2017; 51(12):969–977.
204. Mez J, Daneshvar DH, Kiernan PT, et al. Clinicopathological evaluation of chronic traumatic encephalopathy in players of American Football. *JAMA*. 2017; 318(4):360–370.
205. Belanger HG, Vanderploeg RD, McAllister T. Subconcussive blows to the head: a formative review of short-term clinical outcomes. *J Head Trauma Rehabil*. 2016; 31(3):159–166.
206. Alosco ML, Barr WB, Banks SJ, et al. Neuropsychological test performance of former American football players. *Alzheimers Res Ther*. 2023; 15(1):1.
207. Buckland ME, Sy J, Szentmariay I, et al. Chronic traumatic encephalopathy in two former Australian National Rugby League players. *Acta Neuropathol Commun*. 2019; 7(1):97.
208. Bieniek KF, Cairns NJ, Crary JF, et al. The second NINDS/NIBIB consensus meeting to define neuropathological criteria for the diagnosis of chronic traumatic encephalopathy. *J Neuropathol Exp Neurol*. 2021; 80(3):210–219.
209. Pullman MY, Dickstein DL, DeKosky ST, Gandy S. Antemortem biomarker support for a diagnosis of clinically probable chronic traumatic encephalopathy. *Mol Psychiatry*. 2017; 22(5):638–639.
210. Katz DI, Bernick C, Dodick DW, et al. National Institute of Neurological Disorders and Stroke Consensus Diagnostic Criteria for Traumatic Encephalopathy Syndrome. *Neurology*. 2021; 96(18):848–863.
211. Cherry JD, Babcock KJ, Goldstein LE. Repetitive Head Trauma Induces Chronic Traumatic Encephalopathy by Multiple Mechanisms. *Semin Neurol*. 2020; 40(4):430–438.
212. Kiernan PT, Montenegro PH, Solomon TM, McKee AC. Chronic traumatic encephalopathy: a neurodegenerative consequence of repetitive traumatic brain injury. *Semin Neurol*. 2015; 35(1):20–28.
213. Schwab N, Hazrati LN. Assessing the limitations and biases in the current understanding of chronic traumatic encephalopathy. *J Alzheimers Dis*. 2018; 64(4):1067–1076.
214. Buckland ME, Affleck AJ, Pearce AJ, Suter CM. Chronic traumatic encephalopathy as a preventable environmental disease. *Front Neurol*. 2022; 13:880905.
215. Nowinski CJ, Bureau SC, Buckland ME, et al. Applying the Bradford Hill criteria for causation to repetitive head impacts and chronic traumatic encephalopathy. *Front Neurol*. 2022; 13:938163.
216. Hill AB. Observation and experiment. *N Engl J Med*. 1953; 248(24):995–1001.

217. Hill AB. The environment and disease: association or causation? *Proc R Soc Med*. 1965; 58(5):295–300.
218. McCann H, Bahar AY, Burkhardt K, et al. Prevalence of chronic traumatic encephalopathy in the Sydney Brain Bank. *Brain Commun*. 2022; 4(4).
219. Postupna N, Rose SE, Gibbons LE, et al. The delayed neuropathological consequences of traumatic brain injury in a community-based sample. *Front Neurol*. 2021; 12:624696.
220. Forrest SL, Kril JJ, Wagner S, et al. Chronic traumatic encephalopathy (CTE) is absent From a European community-based aging cohort while cortical aging-related tau astrogliopathy (ARTAG) is highly prevalent. *J Neuropathol Exp Neurol*. 2019; 78(5):398–405.
221. Koerte IK, Lin AP, Willems A, et al. A review of neuroimaging findings in repetitive brain trauma. *Brain Pathol*. 2015; 25(3):318–349.
222. Gardner RC, Yaffe K. Epidemiology of mild traumatic brain injury and neurodegenerative disease. *Mol Cell Neurosci*. 2015; 66(Pt B):75–80.
223. Dougan BK, Horswill MS, Geffen GM. Athletes' age, sex, and years of education moderate the acute neuropsychological impact of sports-related concussion: a meta-analysis. *J Int Neuropsychol Soc*. 2014; 20(1):64–80.
224. Bigler ED, Stern Y. Traumatic brain injury and reserve. *Handb Clin Neurol*. 2015; 128:691–710.
225. Satz P, Cole MA, Hardy DJ, Rassovsky Y. Brain and cognitive reserve: mediator (s) and construct validity, a critique. *J Clin Exp Neuropsychol*. 2011; 33(1):121–130.
226. Stern Y. What is cognitive reserve? Theory and research application of the reserve concept. *J Int Neuropsychol Soc*. 2002; 8(3):448–460.
227. Jordan BD. Chronic traumatic brain injury associated with boxing. *Semin Neurol*. 2000;20(2):179–85.
228. Guterman A, Smith RW. Neurological sequelae of boxing. *Sports Med*. 1987; 4(3):194–210.
229. McKee AC, Cantu RC, Nowinski CJ, et al. Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J Neuropathol Exp Neurol*. 2009; 68(7):709–735.
230. McKee AC, Stern RA, Nowinski CJ, et al. The spectrum of disease in chronic traumatic encephalopathy. *Brain*. 2013; 136(Pt 1):43–64.
231. De Beaumont L, Theoret H, Mongeon D, et al. Brain function decline in healthy retired athletes who sustained their last sports concussion in early adulthood. *Brain*. 2009; 132(3):695–708.
232. Vagnozzi R, Signoretti S, Tavazzi B, Floris R, Ludovici A, Marziali S, Tarascio G, Amorini AM, Di Pietro V, Delfini R, Lazzarino G. Temporal window of metabolic brain vulnerability to concussion: a pilot 1H-magnetic resonance spectroscopic study in concussed athletes-part III. *Neurosurgery*. 2008 Jun;62(6):1286–95.
233. Orchard JW, Orchard JJ, Semsarian C, La Gerche A, Driscoll T. Reduced death rates of elite Australian Rules footballers compared to age-matched general population. *J Sci Med Sport*. 2022; 25(9):710–714.
234. Baron SL, Hein MJ, Lehman E, Gersic CM. Body mass index, playing position, race, and the cardiovascular mortality of retired professional football players. *Am J Cardiol*. 2012; 109(6):889–896.
235. Webner D, Iverson GL. Suicide in professional American football players in the past 95 years. *Brain Inj*. 2016; 30(13–14):1718–1721.
236. Iverson GL. Chronic traumatic encephalopathy and risk of suicide in former athletes. *Br J Sports Med*. 2014; 48(2):162–164.
237. Iverson GL. Suicide and chronic traumatic encephalopathy. *J Neuropsychiatry Clin Neurosci*. 2015; 28(1):9–16.
238. Wright DK, Gardner AJ, Wojtowicz M, et al. White matter abnormalities in retired professional rugby league players with a history of concussion. *J Neurotrauma*. 2021; 38(8):983–988.
239. Guell X, Arnold Anteraper S, Gardner AJ, et al. Functional connectivity changes in retired rugby league players: a data-driven functional magnetic resonance imaging study. *J Neurotrauma*. 2020; 37(16):1788–1796.
240. Gardner AJ, Iverson GL, Wojtowicz M, et al. MR Spectroscopy findings in retired professional rugby league players. *Int J Sports Med*. 2017; 38(3):241–252.
241. Iverson GL, Van Patten R, Terry DP, Levi CR, Gardner AJ. Predictors and correlates of depression in retired elite level rugby league players. *Front Neurol*. 2021; 12.
242. Van Patten R, Iverson GL, Terry DP, Levi CR, Gardner AJ. Predictors and correlates of perceived cognitive decline in retired professional rugby league players. *Front Neurol*. 2021; 12.
243. Stanwell P, Iverson GL, Van Patten R, Castellani RJ, McCrory P, Gardner AJ. Examining for cavum septum pellucidum and ventricular enlargement in retired elite-level rugby league players. *Front Neurol*. 2022; 13.
244. Beam CR, Kaneshiro C, Jang JY, Reynolds CA, Pedersen NL, Gatz M. Differences between women and men in incidence rates of dementia and Alzheimer's disease. *J Alzheimers Dis*. 2018; 64(4):1077–1083.

245. Johnson VE, Stewart W, Smith DH. Traumatic brain injury and amyloid- β pathology: a link to Alzheimer's disease? *Nat Rev Neurosci.* 2010; 11(5):361–370.
246. Angum F, Khan T, Kaler J, Siddiqui L, Hussain A. The prevalence of autoimmune disorders in women: a narrative review. *Cureus.* 2020; 12(5):e8094–e8094.
247. Eimer WA, Vijaya Kumar DK, Navalpur Shanmugam NK, et al. Alzheimer's disease-associated β -Amyloid is rapidly seeded by Herpesviridae to protect against brain infection. *Neuron.* 2018; 99(1):56–63.e53.
248. Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol.* 2016; 16(10):626–638.
249. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet.* 2020; 396(10248):413–446.
250. Stern Y, Arenaza-Urquijo EM, Bartrés-Faz D, et al. Whitepaper: Defining and investigating cognitive reserve, brain reserve, and brain maintenance. *Alzheimers Dement.* 2020; 16(9):1305–1311.
251. Buckland ME, Affleck AJ, Pearce AJ, Suter CM. Chronic Traumatic Encephalopathy as a Preventable Environmental Disease. *Front Neurol.* 2022; 13.
252. Selvackadunco S, Langford K, Shah Z, et al. Comparison of clinical and neuropathological diagnoses of neurodegenerative diseases in two centres from the Brains for Dementia Research (BDR) cohort. *J Neural Transm.* 2019; 126(3):327–337.
253. SantaCruz KS, Sonnen JA, Pezhouh MK, Desrosiers MF, Nelson PT, Tyas SL. Alzheimer disease pathology in subjects without dementia in 2 studies of aging: the Nun Study and the Adult Changes in Thought Study. *J Neuropathol Exp Neurol.* 2011; 70(10):832–840.
254. Stern Y, Gurland B, Tatemichi TK, Tang MX, Wilder D, Mayeux R. Influence of education and occupation on the incidence of Alzheimer's disease. *JAMA.* 1994; 271(13):1004–1010.
255. Brayne C, Ince PG, Keage HA, et al. Education, the brain and dementia: neuroprotection or compensation? *Brain.* 2010; 133(Pt 8):2210–2216.
256. Del Ser T, Hachinski V, Merskey H, Munoz DG. An autopsy-verified study of the effect of education on degenerative dementia. *Brain.* 1999; 122 (Pt 12):2309–2319.
257. Farfel JM, Nitrini R, Suemoto CK, et al. Very low levels of education and cognitive reserve: a clinicopathologic study. *Neurology.* 2013; 81(7):650–657.
258. Fratiglioni L, Paillard-Borg S, Winblad B. An active and socially integrated lifestyle in late life might protect against dementia. *Lancet Neurol.* 2004; 3(6):343–353.
259. Scarmeas N, Levy G, Tang MX, Manly J, Stern Y. Influence of leisure activity on the incidence of Alzheimer's disease. *Neurology.* 2001; 57(12):2236–2242.
260. Wang HX, Xu W, Pei JJ. Leisure activities, cognition and dementia. *Biochim Biophys Acta.* 2012; 1822(3):482–491.
261. Gardner AJ, Kay-Lambkin F, Shultz SR, Iverson GL. Level of knowledge and attitude towards sport-related concussion among the general? public. *Br J Sports Med.* 2017; 51(11):A68–A68.
262. Pearce AJ, Young JA, Parrington L, Aimers N. Do as I say: contradicting beliefs and attitudes towards sports concussion in Australia. *J Sports Sci.* 2017; 35(19):1911–1919.
263. White PE, Register-Mihalik J, Donaldson A, Sullivan SJ, Finch CF. Concussion guideline implementation perceptions and experiences among parents of community-level Australian Football junior players. *BMJ Open Sport Exerc Med.* 2017; 3(1):e000215.
264. Emery CA, Black AM, Kolstad A, et al. What strategies can be used to effectively reduce the risk of concussion in sport? A systematic review. *Br J Sports Med.* 2017; 51(12):978–984.
265. Taylor ME, Sanner JE. The relationship between concussion knowledge and the high school athlete's intention to report traumatic brain injury symptoms: a systematic review of the literature. *J Sch Nurs.* 2017; 33(1):73–81.
266. Clacy A, Goode N, Sharman R, Lovell GP, Salmon P. A systems approach to understanding the identification and treatment of sport-related concussion in community rugby union. *Appl Ergon.* 2019; 80:256–264.
267. White PE, Donaldson A, Sullivan SJ, Newton J, Finch CF. Australian Football League concussion guidelines: what do community players think? *BMJ Open Sport Exerc Med.* 2016; 2(1):e000169.

APPENDIX A — PHYSIOTHERAPY GUIDED REHABILITATION OF CONCUSSION

The following table outlines the five systems that should be considered when designing a rehabilitation plan for an athlete recovering from concussion and is based on the graded return to sport framework (GRTSF) for **advanced care settings**. The systems are closely interconnected and care must be taken to avoid over stimulation, particularly early in the GRTSF. Failure to progress through the graded return to sport as expected requires referral to appropriate member of the clinical management team (or CMT) for subsequent review. Athletes need to be assessed individually as the requirements for every sport and athlete are different and no two concussions present the same. The below is an example and should not be used as a recipe. Some considerations are included at each stage that can challenge/be useful for practitioners and coaches.

Athletes 18 years and under can utilise this guide but MUST follow the GRTSF for community and youth, specifically the 14 days symptom free (at rest) and 21 days minimum before returning to competitive contact.

Autonomic	Cervical	Vestibular	Visual	Cognitive
Activities of daily living	Full pain-free range of motion (active and passive) Physiotherapy assessment will guide rehabilitation but can involve passive, active and active-assisted range of motion exercises	Activities of daily living	Activities of daily living	Activities of daily living
Steady-state exercise Up to 60% HRmax with minimal movement of head e.g. moderate walk over flat ground or stationary bike seated only	Address referred pain and headaches Physiotherapist-led treatment as appropriate for each individual. May include manual therapy, exercise therapy		Gradual resumption of school or work e.g. depending on symptoms begin with several hours or part days at work or school <u>Consider:</u> Decrease work of visual system by sitting at the front of classrooms, scheduling short periods of time at the computer. Avoid watching fast paced or complex visuals such as training sessions, videos, movies, etc	
Healthcare practitioner review recommended at day 3-4 to include SCOAT6				
Continue steady-state exercise (as above)	Deep neck flexion exercises initiated Motor control and stability exercises can be prescribed in seated, supine or prone positions depending on individual requirements. <u>Consider:</u> Sport-specific positions for head/neck such as: > 4-point kneeling for rugby, cycling > prone holds with rotation for swimming, gymnastics > seated for waterpolo, artistic swimming > standing multidirectional for AFL	Continue activities of daily living (as above)	Continue gradual resumption of school or work (as above)	Continue gradual resumption of school or work Progressions will naturally increase as rehabilitation in other areas progresses. If complications and impairments in cognitive function are recognised during rehabilitation referral to a HCP* is recommended for assessment. If there are known impairments in cognitive function post-concussion close communication with the managing doctor is highly recommended for an integrated management approach and best outcomes to occur
			Hand-eye coordination with minimal head movements e.g. > Ball drills against wall or with partner from in front only > Steady running at 50-60% max speed > Swimming with kickboard and snorkel <u>Consider:</u> Speed, height, type of ball (if using) and predictability of activity performed. Start in quiet environment with bland visual background to avoid exacerbation of symptoms due to visual and cognitive loads	
Progress steady-state exercise Maintain single plane of movement, e.g. straight line running, etc. Incremental increase in heart rate up to 80% HRmax e.g. 5min increments @50%, 60%, 65%, 70% Hrmx e.g. start with stationary seated bike or walking and progress to sport-specific mediums such as: > Running (straight line only) > Swimming (minimal to no rotation, no tumbling at ends of pool) > Cycling return to road riding (bike paths or quiet roads) <u>Consider:</u> Concussive symptoms may be exacerbated by environments that challenge the visual or vestibular system. These can include: noise, lights, movements in athlete's visual background, team/squad training, uneven surfaces, etc.	Cervical sensorimotor exercises e.g. joint position training seated with eyes open <u>Consider:</u> Progress difficulty with eyes closed or standing on firm and soft surfaces. Visual backgrounds, sport-specific environments can increase complexity such as: > Seated on exercise ball > Eggbeater position on plinth and progressing to pool > 4-point kneeling > Prone	Controlled non-reactive hand-eye coordination with head movements e.g. > Ball drills against wall or with partner from various directions to encourage head rotations, up and downs, etc. > Increase running pace to add greater visual input, run around track > Swimming strokes with usual movement patterns (no tumbling) <u>Consider:</u> Complex visual backgrounds, busy environments such as sideline of team/squad training, crowd noises, lights can challenge visual and cognitive system but may also exacerbate symptoms		Introduce dual-tasks e.g. > Count numbers out loud whilst doing ball skills > Complete multiplications on demand whilst completing steady state exercise, such as stationary bike or walking

Autonomic	Cervical	Vestibular	Visual	Cognitive
Do any activities bring on or exacerbate symptoms? Y N				
Can athlete complete 1-minute of sport-specific skills with head movement without ANY symptoms? Y N				
Continue progressing steady-state exercise (as above)	Introduce resistance training Build strength loads back to squad/individual requirements. Include specific cervical flexor and extensor strengthening. <u>Consider:</u> Re-integration with squad/team during S&C sessions to challenge visual, vestibular, and cognitive loads as appropriate	Continue controlled non-reactive hand-eye coordination with head movements (as above)	See above for details	
		More complex hand-eye (or foot-eye) coordination involving head and body movements e.g. > Ball drills involving 90degree rotations progressing to 360deg > Introduce small volumes of tumbling at ends of pool and diving from blocks > Starting from blocks and reacting to start signal > Shooting practice from greater distances (waterpolo, soccer, AFL, archery, shooting, basketball, etc.) <u>Consider:</u> Volume of skills being introduced. Sport specific coordination skills, e.g. groundballs, high balls, cycling with visual input e.g. simulator and increase complexity by performing tasks in busy training environments and progressing pace of tasks.		
Interval training > increase run speeds to 90% with straight-line run-throughs, flying 60s, etc. > swimming interval sets <u>Consider:</u> Ensure time allocated between sets for recovery and any potential symptom provocation	Continue resistance training (as above)	Hand-eye (or foot/eye) coordination whilst moving e.g. > walk-through ball skills such as walking dribbling, partner passing whilst walking > progress skills involving rotations, twisting, precision, e.g. gymnastics, diving, artistic swimming, archery, shooting <u>Consider:</u> Environment of activities and increase complexity with multiple balls and/or players, performing multiple skills consecutively, volume, noise, lights, etc		
Agility and multi-directional activity e.g. > incorporate planned or athlete-led directional changes into running > pool sessions with unrestricted tumbling > progressively return to busier roads on bike <u>Consider:</u> Progress to reactionary change of directions to increase cognitive load. Use speed, volume, sport specific skills to increase challenge		Controlled sport-specific activities e.g. Controlled team non-contact or high-risk training activities such as: > kicking or hand-balling drills (No match simulation drills) > 5v0 drills <u>Consider:</u> Volume if all other parameters are back to full training capacity. Number of athletes, noise, external and internal stress may increase cognitive and visual load		
Build training volumes and sport-specific requirements Incremental increase to meet usual training volumes for squad/individual <u>Consider:</u> All systems being rehabilitated simultaneously		Reactive sport-specific activities e.g. Uncontrolled non-contact training or physiotherapist-led reactive drills. involving: > Kick, chase, marking skills > Rebounding drills > Uncontrolled terrains <u>Consider:</u> Vary reactionary component with location, timing or skill required. Increase complexity with multiple balls and/or players, competition against teammates, noise, lights, etc		
Healthcare practitioner review for clearance to return to contact and high-risk activities When symptom-free for at least 10 days				
Full Training				
Competition or Match Simulation				
Match or Competition Play				



Australian Government
Australian Sports Commission

ALS.gov.au



@theAIS #theAIS